

# MA5204 Commutative and Homological Algebra

Thang Pang Ern

## Contents

---

<b>1</b>	<b>Recap of Ring Theory and Module Theory</b>	<b>2</b>
1.1	Ring Theory . . . . .	2
1.2	Module Theory . . . . .	14
<b>2</b>	<b>Basic Commutative Algebra</b>	<b>16</b>
2.1	Exact Sequences of Modules . . . . .	16
2.2	Localization . . . . .	20
2.3	Tensor Products . . . . .	25
<b>3</b>	<b>Some Classes of Rings</b>	<b>32</b>
3.1	Noetherian Rings . . . . .	32
3.2	A Preview of Dimension Theory . . . . .	33
3.3	Integral Dependence and Integral Rings . . . . .	34
3.4	Completions . . . . .	39
<b>4</b>	<b>Introduction to Homology</b>	<b>42</b>

## References

---

- 1 Atiyah, M., Macdonald, I. (1994). *Introduction to Commutative Algebra*. CRC Press.
- 2 Matsumura, H. (1986). *Commutative Ring Theory*. Cambridge University Press.
- 3 Eisenbud, D. (1995). *Commutative Algebra*. Springer, Berlin.

# Chapter 1

## Recap of Ring Theory and Module Theory

---

### 1.1 Ring Theory

**Definition 1.1 (ring).** A ring  $R$  is a set with distinct elements  $1, 0 \in R$  equipped with two binary maps which are multiplication and addition respectively.

$$R \times R \rightarrow R \text{ where } (r, r') \mapsto rr' \quad \text{and} \quad R \times R \times R \text{ where } (r, r') \mapsto r + r'.$$

The following conditions are satisfied:

(i)  $(R, +, 0)$  is an Abelian group, i.e. for all  $r, r' \in R$ ,

$$r + r' = r' + r \quad \text{and} \quad 0 + r = r = r + 0$$

(ii) Distributivity and associativity holds, i.e. for all  $r, s, s_1, s_2, t \in R$ ,

$$r(s_1 + s_2) = rs_1 + rs_2 \quad \text{and} \quad r(st) = (rs)t$$

(iii) Existence of multiplicative identity, i.e.  $1r = r1 = r$  for all  $r \in R$

We say that  $R$  is an associative ring with unity.

**Definition 1.2 (commutative ring).** If we further assume that  $rs = sr$  for all  $r, s \in R$  in Definition 1.1, we obtain a commutative ring with unity.

**Remark 1.1.** In this course, we take rings to be *commutative rings with unity*.

**Definition 1.3 (unit).** Let  $x \in R$ . If

there exists  $y \in R$  such that  $xy = 1$  then  $x$  is a unit.

Here,  $y = 1/x$ .

**Proposition 1.1.** The set of units of  $R$ , denoted by  $R^\times$ , forms an Abelian group under  $\times$ .

**Definition 1.4 (field).** A ring  $R$  is a field if  $R^\times = R \setminus \{0\}$ .

**Definition 1.5 (ring homomorphism).** A ring homomorphism  $\varphi : R \rightarrow S$  is a map of sets such that

- (i)  $\varphi(0_R) = 0_S$
- (ii)  $\varphi(1_R) = 1_S$
- (iii)  $\varphi(r + r') = \varphi(r) + \varphi(r')$
- (iv)  $\varphi(rr') = \varphi(r)\varphi(r')$

**Definition 1.6 (ideal).** Let  $R$  be a ring. An ideal of  $R$  is a subset  $I \subseteq R$  such that

(i)  $I \leq (R, 0, +)$ , i.e.

$$0 \in I \quad \text{and} \quad \text{for all } i_1, i_2 \in I \text{ we have } i_1 + i_2 \in I$$

(ii) For all  $r \in R$  and  $i \in I$ , we have  $ri \in I$

**Example 1.1 (integer multiples).** For any fixed integer  $n \in \mathbb{Z}$ ,

$$n\mathbb{Z} = \{\text{all multiples of } n\} \subseteq \mathbb{Z} \quad \text{is an ideal.}$$

**Example 1.2.** More generally, given any  $x \in R$ , the subset

$$(x) = \{\text{all elements in } R \text{ of the form } xr : r \in R\} \subseteq R \quad \text{is an ideal.}$$

**Proposition 1.2.** If  $I \subseteq R$  is an ideal, then the set

$$R/I = \text{quotient of } R \text{ by } I \text{ as Abelian groups} = \text{the set of cosets } r + I \subseteq R$$

naturally has a ring structure.

*Proof.* Let  $r_1, r_2 \in R$ . We have

$$(r_1 + I) + (r_2 + I) = r_1 + r_2 + I \quad \text{and} \quad (r_1 + I)(r_2 + I) = r_1 r_2 + I.$$

Also,  $1 = 1_R + I$  and  $0 = 0_R + I$ . Note that by construction, there exists a natural surjective ring homomorphism  $R \rightarrow R/I$ , i.e. any surjective ring homomorphism  $f : R \rightarrow S$  arises from such a construction if we set  $I = f^{-1}(0)$ , so  $S \cong R/I$ .  $\square$

**Example 1.3.** Let  $R = \mathbb{Z}$  and  $I = (n)$ . Then,

$$R/I = \mathbb{Z}/(n) = \{0, 1, \dots, n-1\} \quad \text{which is precisely the integers modulo } n.$$

A simple fact from MA1100 states that that  $\mathbb{Z}/(n)$  is a field if and only if  $n$  is some prime  $p$ .

**Definition 1.7 (integral domain).** A ring  $R$  is a integral domain if

$$\text{for all } x, y \in R, \text{ we have } xy = 0 \quad \text{implies} \quad x = 0 \text{ or } y = 0.$$

**Definition 1.8 (prime ideal).** Let  $A$  be a ring. An ideal  $I \subseteq A$  is prime if

$$\text{for all } x, y \in A, \text{ we have } xy \in I \quad \text{implies} \quad x \in I \text{ or } y \in I.$$

**Proposition 1.3.** Let  $A$  be a ring. Given any  $I \subseteq A$ ,

$$A/I \text{ is an integral domain} \quad \text{if and only if} \quad I \text{ is a prime ideal.}$$

*Proof.* We only prove the reverse direction. The proof of the forward direction is similar. Anyway, given  $x, y \in A$  for some ring  $A$ , suppose  $I$  is a prime ideal. Say  $\bar{x} \cdot \bar{y} = 0$ . This holds if and only if  $xy \in I$ . Equivalently,  $x \in I$  or  $y \in I$ , i.e.  $\bar{x} = 0$  or  $\bar{y} = 0$ . As such,  $A/I$  is an integral domain.  $\square$

**Definition 1.9 (maximal ideal).** An ideal  $I \subset A$  (proper subset inclusion) is maximal if there does not exist any ideals  $I \subset J \subset A$ .

**Proposition 1.4.** Let  $A$  be a ring. Then,

an ideal  $I \subset A$  is maximal if and only if  $A/I$  is a field.

*Proof.* Note that given any ring homomorphism  $\varphi : A \twoheadrightarrow A/I$  in  $A$ , there is a natural inclusion-preserving bijection between

$$\{\text{ideals } I \subseteq J \subseteq A\} \quad \text{and} \quad \{\text{ideals } \bar{J} \subseteq A/I\}.$$

The map is given by  $J \mapsto J/I = \bar{J}$  such that  $\bar{J} \mapsto \varphi^{-1}(\bar{J})$  since  $\varphi$  is bijective, hence invertible.

Now, consider the following chain of implications:

$J \subset A$  is maximal if and only if the only ideals of  $A/I$  are  $A/I$  and  $(0)$   
 if and only if any  $0 \neq x \in A/I$  satisfies  $(x) = A/I$   
 if and only if any  $0 \neq x \in A/I$  is a unit  
 if and only if  $A/I$  is a field

The result follows. □

**Proposition 1.5.** Any non-zero ring  $A$  has a maximal ideal.

*Proof.* Recall Zorn's lemma which states that if  $S \neq \emptyset$  is a partially ordered set such that any chain in  $S$  admits an upper bound, then  $S$  has a maximal element. Recall that a chain  $C$  is a subset of  $S$  such that

$$\text{for all } x, y \in S \quad \text{we have} \quad x \leq y \text{ or } y \leq x.$$

Now, fix a non-zero ring  $A$ . Let  $S$  denote the set of proper ideals  $I \subset A$  with the inclusion being the partial order relation. Note that  $S \neq \emptyset$  since  $(0) \in S$ . Next, if  $C \subseteq S$  is a chain, then

$$\bigcup_{s \in C} I_s \quad \text{is a proper ideal.}$$

Thus, the aforementioned union is contained in  $S$  and is an upper bound for the chain  $C$ .

As such, Zorn's lemma applies so  $S$  has a maximal element if and only if  $A$  has a maximal ideal. □

**Corollary 1.1.** For any ring  $A$ ,

any proper ideal  $I \subset A$  is contained in some maximal ideal.

*Proof.* Suppose  $I$  is a proper ideal of  $A$ . Then,  $A/I \neq 0$ , which implies that there exists a maximal ideal  $\mathfrak{m}$  properly contained in  $A/I$ . So, the preimage of  $\mathfrak{m}$  in  $A$  is maximal and contains  $I$ . □

**Definition 1.10 (nilpotent element).** Let  $A$  be a ring. An element  $x \in A$  is nilpotent if

$$\text{there exists } n \in \mathbb{N} \text{ such that } x^n = 0.$$

**Example 1.4.** 0 is always nilpotent.

**Example 1.5.**  $2 \in \mathbb{Z}/(4)$  is non-zero and nilpotent.

**Example 1.6 (Atiyah and Macdonald p. 10 Question 2).** Let  $A$  be a ring and let  $A[x]$  be the ring of polynomials in an indeterminate  $x$ , with coefficients in  $A$ . Let

$$f = a_0 + a_1x + \dots + a_nx^n \in A[x].$$

Prove that:

- (i)  $f$  is a unit in  $A[x]$  if and only if  $a_0$  is a unit in  $A$  and  $a_1, \dots, a_n$  are nilpotent  
*Hint:* If  $b_0 + b_1x + \dots + b_mx^m$  is the inverse of  $f$ , prove by induction on  $r$  that  $a_n^{r+1}b_{m-r} = 0$ . Hence show that  $a_n$  is nilpotent, and then use the following fact: if  $x$  a nilpotent element of a ring  $A$ , then  $1 + x$  is a unit of  $A$ , for which it follows that the sum of a nilpotent element and a unit is a unit.
- (ii)  $f$  is nilpotent if and only if  $a_0, a_1, \dots, a_n$  are nilpotent
- (iii)  $f$  is a zero-divisor if and only if there exists  $a \neq 0$  in  $A$  such that  $af = 0$   
*Hint:* Choose a polynomial  $g = b_0 + b_1x + \dots + b_mx^m$  of least degree  $m$  such that  $fg = 0$ . Then  $a_nb_m = 0$ , hence  $a_ng = 0$  (because  $a_n$  annihilates  $f$  and has degree  $< m$ ). Now show by induction that  $a_n^r g = 0$  ( $0 \leq r \leq n$ ).
- (iv)  $f$  is said to be primitive if  $(a_0, a_1, \dots, a_n) = (1)$ . Prove that if  $f, g \in A[x]$ , then

$$fg \text{ is primitive if and only if } f \text{ and } g \text{ are primitive.}$$

*Solution.*

- (i) We only prove the forward direction. The proof of the reverse direction follows from the hint (which is actually Question 1 of the same exercise set) and (ii) of this exercise. Suppose  $f$  is a unit in  $A[x]$ . Let  $g = b_0 + b_1x + \dots + b_mx^m$  be the inverse of  $f$ . Then,

$$fg = (a_0 + a_1x + \dots + a_nx^n)(b_0 + b_1x + \dots + b_mx^m)$$

Since the constant term must be 1, then  $a_0b_0 = 1$ , so  $a_0$  is a unit in  $A$ . Recall the convolution formula that

$$fg = c_0 + c_1x + \dots + c_kx^k,$$

where  $c_0 = a_0b_0$  (discussed earlier),

$$c_1 = a_0b_1 + a_1b_0 = 0$$

$$c_2 = a_0b_2 + a_1b_1 + a_2b_0 = 0$$

and so on. One can deduce that  $a_1, \dots, a_n$  are nilpotent.

- (ii) For the forward direction, suppose  $f$  is nilpotent. Then, one can apply induction to  $n$  to show that all of its coefficients are nilpotent. To demonstrate this, note that the  $n = 1$  case is trivial. For the general case, the leading coefficient will be  $a_n^k$  for some  $k \in \mathbb{N}$ , so  $a_n$  is nilpotent. By the inductive hypothesis,  $a_0, \dots, a_{n-1}$  are nilpotent as well.

For the reverse direction, if  $a_0, \dots, a_n$  are nilpotent, define  $d \in \mathbb{N}$  such that

$$a_i^d = 0 \quad \text{for all } 0 \leq i \leq n.$$

In other words,  $d$  is the sum of the orders of all the orders of the coefficients. As such,  $f^d = 0$ .

- (iii) For the forward direction, suppose  $f$  is a zero divisor. Then, let  $g$  be a polynomial of minimal order such that  $fg = 0$ . Suppose  $g = b_0 + b_1x + \dots + b_mx^m$  such that  $\deg g > 0$ . Then,  $a_nb_m = 0$ , i.e.  $a_n g$  annihilates  $f$  but  $\deg(a_n g) < m$ , which is a contradiction. As such,

$$\deg g = 0 \quad \text{or in other words} \quad \text{there exists } a \in A \text{ such that } af = 0.$$

The reverse direction follows by the definition of a zero-divisor (recall MA3201).

- (iv) The reverse direction is essentially Gauss' lemma (MA3201); for the forward direction, if  $fg$  is primitive, then  $(c_0, \dots, c_{n+m}) = (1)$ , where the  $c_i$ 's are the coefficients of  $fg$ . This means that  $\gcd(c_0, \dots, c_{n+m}) = 1$ , or equivalently, there does not exist  $d > 1$  which divides all the  $c_i$ 's.

Suppose on the contrary that neither  $f$  nor  $g$  is primitive. Then, say  $\gcd(a_0, \dots, a_n) > 1$ . Then, because of the convolution formula

$$c_k = \sum_{i+j=k} a_i b_j \quad (\text{look at the dependence between } a_i \text{ and } c_k),$$

it forces the existence of some  $d > 1$  which divides all the  $c_i$ 's, leading to a contradiction! □

**Proposition 1.6 (nilradical).** The set of nilpotent elements in any ring  $A$  is an ideal. We call this the nilradical of  $A$  which is denoted by  $\mathfrak{N}_A$ .

*Proof.* Suppose  $x \in A$  is nilpotent, i.e.

$$\text{there exists } n \in \mathbb{N} \quad \text{such that} \quad x^n = 0.$$

Then, for any  $r \in A$ , we have

$$(rx)^n = r^n x^n = r^n \cdot 0 = 0.$$

For compatibility regarding addition, suppose  $x, y \in A$  are nilpotent. Then,

$$\text{there exist } n, m \in \mathbb{N} \quad \text{such that} \quad x^n = 0 \text{ and } y^m = 0.$$

We use the binomial theorem to obtain

$$(x + y)^{n+m} = x^{n+m} + \binom{n+m}{1} x^{n+m-1} y + \dots + \binom{n+m}{m} x^n y^m + \dots + \binom{n+m}{n+m-1} x y^{n+m-1} + y^{n+m}$$

which is 0 (*not surprising anyway*). □

**Definition 1.11 (reduced ring).** A ring  $A$  is reduced if it contains no non-zero nilpotent elements.

**Example 1.7.** A nice observation: for  $n \neq 0$ ,

$$\mathbb{Z}/(n) \text{ is reduced} \quad \text{if and only if} \quad n \text{ is squarefree.}$$

**Proposition 1.7.** For any non-zero  $A$ , we have

$$\mathfrak{N}_A = \bigcap_{\mathfrak{p} \subset A} \mathfrak{p},$$

where  $\mathfrak{p}$  denotes a prime ideal of  $A$ .

*Proof.* We first prove the forward inclusion. Suppose  $x \in A$  is nilpotent. Then,  $\bar{x} \in A/\mathfrak{p}$  is nilpotent, so  $\bar{x} = 0$  in  $A/\mathfrak{p}$  since  $A/\mathfrak{p}$  is an integral domain. As such,  $x \in \mathfrak{p}$  for all  $\mathfrak{p} \subset A$ .

For the reverse direction, fix  $x \notin \mathfrak{N}_A$ . We wish to find a prime ideal  $\mathfrak{p}$  such that  $x \notin \mathfrak{p}$ . Let

$$\Sigma = \{I \subset A : x^n \notin I \text{ for all } n \in \mathbb{N}\}.$$

Then,  $\Sigma \neq \emptyset$  as  $(0) \in \Sigma$  by assumption on  $x$ . By applying the same argument as before, any chain in  $\Sigma$  has an upper bound. By Zorn's lemma,  $\Sigma$  has a maximal element  $\mathfrak{p}$ . It suffices to show that  $\mathfrak{p}$  is a prime ideal. Suppose  $y, z \in A \setminus \mathfrak{p}$ . We wish to show that  $yz \notin \mathfrak{p}$ . Note that

$$\mathfrak{p} \subset (\mathfrak{p}, y) \quad \text{and} \quad \mathfrak{p} \subset (\mathfrak{p}, z).$$

These imply the following respectively: there exist  $n, m \in \mathbb{N}$  such that  $x^n \in (\mathfrak{p}, y)$  and  $x^m \in (\mathfrak{p}, z)$ . So,

$$x^n = p_1 + yr_1 \quad \text{and} \quad x^m = p_2 + zr_2 \quad \text{for } p_1, p_2 \in \mathfrak{p} \text{ and } r_1, r_2 \in A.$$

Multiplying both elements, we obtain

$$\mathfrak{p} \not\supseteq x^{n+m} = p_1p_2 + p_1zr_2 + p_2yr_1 + yzr_1r_2 \in yzr_1r_2 + \mathfrak{p}.$$

Hence,  $yzr_1r_2 \notin \mathfrak{p}$  and the result follows.  $\square$

**Example 1.8** (Atiyah and Macdonald p. 11 Question 8). Let  $A$  be a ring  $\neq 0$ . Show that the set of prime ideals of  $A$  has a minimal element with respect to inclusion.

*Solution.* Note that every descending chain of prime ideals  $\mathfrak{p}$  has a lower bound, which is their intersection. By Zorn's lemma, the set of prime ideals of  $A$  has at least one minimal element.  $\square$

**Remark 1.2.** Similar to Example 1.13, the set of prime ideals of  $A$  in Example 1.8 is actually called the prime spectrum of  $A$  or  $\text{Spec}(A)$ .

Given two ideals  $I, J \subseteq R$ , we can construct some *new* ideals (Proposition 1.8).

**Proposition 1.8 (constructing new ideals).** Fix a ring  $R$ . Suppose we are given ideals  $I, J \subseteq R$ . Then, the following are also ideals of  $R$ :

- (i)  $I \cap J$
- (ii)  $I + J = \{i + j : i \in I, j \in J\}$
- (iii)  $IJ = \{i_1j_1 + \dots + i_kj_k : i_m \in I, j_n \in J\}$

We have obvious generalisations to ideals  $I_1, \dots, I_n \subseteq R$ .

**Proposition 1.9.** If  $x_1, \dots, x_n \in R$  are given, we call

$$(x_1, \dots, x_n) = (x_1) + \dots + (x_n) \quad \text{the ideal generated by } x_1, \dots, x_n.$$

**Example 1.9.** Let  $R = \mathbb{Z}$ , i.e. the ring of integers and consider the ideals  $I = (n)$  and  $J = (m)$ . Then,

$$\begin{aligned} IJ &= (nm) \\ I+J &= (\gcd(m, n)) \\ I \cap J &= (\text{lcm}(m, n)) \end{aligned}$$

**Proposition 1.10.** Fix a ring  $R$ . Suppose we have ideals  $I, J \subseteq R$ . Then, the following hold:

- (i)  $IJ \subseteq I \cap J \subseteq I + J$
- (ii) In general, we have  $(I+J)(I \cap J) \subseteq IJ$ . In fact, if  $I$  and  $J$  are coprime (that is  $I+J = R$ ), then  $IJ = I \cap J$ .

**Proposition 1.11.** Let  $R$  be a ring. Suppose we have ideals  $I, J \subseteq R$ . Consider the ring multiplication

$$\varphi : R \rightarrow R/I \times R/J.$$

Then, the following hold:

- (i)  $\ker \varphi = I \cap J$
- (ii) If  $I+J = R$ , i.e.  $I$  and  $J$  are coprime, then  $\varphi$  is surjective
- (iii) If  $I$  and  $J$  are coprime, then we have the isomorphism

$$R/IJ \cong R/(I \cap J) \cong R/I \times R/J$$

In fact, the Chinese remainder theorem states that  $R/IJ \cong R/I \times R/J$ .

*Proof.* We will only prove (ii) and (iii) as the proof of (i) is obvious. For (ii), choose  $\bar{x} \in R/I$  and  $\bar{y} \in R/J$ . Since  $I+J = R$ , then we can write

$$1 = i + j \quad \text{for some } i \in I, j \in J.$$

Note that

$$\varphi(i) = (0, 1) \quad \text{and} \quad \varphi(j) = (1, 0).$$

Since  $\varphi$  is a ring homomorphism, then

$$\varphi(jx + iy) = (\bar{x}, \bar{y}) \quad \text{which shows that } \varphi \text{ is surjective.}$$

Moreover,  $\ker \varphi = I \cap J = IJ$ . Thus, the isomorphism in (iii) holds. □

**Proposition 1.12 (extension and contraction).** Suppose  $\varphi : R \rightarrow S$  is a ring homomorphism. We have

$$J \subseteq S \text{ is an ideal} \quad \text{implies} \quad \varphi^{-1}(J) \subseteq R \text{ is an ideal.}$$



This is often called the contraction of  $I$ . However, if  $I \subseteq R$  is an ideal, then  $\varphi(I) \subseteq S$  need not be an ideal. So, we can consider  $\varphi(I)S$  to be the ideal generated by  $\varphi(I)$  (called the extension of  $I$  along  $\varphi$ ), where

$$\varphi(I)S = \{s_1\varphi(i_1) + \dots + s_k\varphi(i_k) : s_m \in S, i_m \in I\}$$

**Example 1.10.** Given the inclusion map  $\varphi : \mathbb{Z} \hookrightarrow \mathbb{Q}$ , the zero ideal is maximal in  $\mathbb{Z}$ , but its pre-image is not maximal in  $\mathbb{Q}$ . Thus the pre-image of a maximal ideal is not necessarily maximal.

**Proposition 1.13.** Let  $\varphi : R \rightarrow S$  be a ring homomorphism. Then,

$$\mathfrak{p} \subseteq S \text{ is a prime ideal} \implies \varphi^{-1}(\mathfrak{p}) \subseteq R \text{ is also a prime ideal.}$$

*Proof.* The following diagram commutes:

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \downarrow & & \downarrow \\ R/\varphi^{-1}(\mathfrak{p}) & \longrightarrow & S/\mathfrak{p} \end{array}$$

and the lower horizontal arrow is injective. That is, we have  $R/\varphi^{-1}(\mathfrak{p}) \hookrightarrow S/\mathfrak{p}$ . Then,  $\mathfrak{p}$  is prime if and only if  $S/\mathfrak{p}$  is an integral domain, and equivalently,  $R/\varphi^{-1}(\mathfrak{p}) \subseteq S/\mathfrak{p}$  is an integral domain. We conclude that  $\varphi^{-1}(\mathfrak{p})$  is a prime ideal.  $\square$

**Definition 1.12 (prime spectrum).** For any ring  $R$ , let  $\text{Spec } R$  denote the set of all prime ideals of  $R$ . That is,

$$\text{Spec } R = \{\mathfrak{p} \subseteq R : \mathfrak{p} \text{ is prime}\}.$$

Proposition 1.13 implies that for any ring homomorphism  $\varphi : R \rightarrow S$ , there exists an induced homomorphism  $\varphi^* : \text{Spec } S \rightarrow \text{Spec } R$ , where  $\varphi^*(\mathfrak{p}) = \varphi^{-1}(\mathfrak{p})$ . We can upgrade  $\text{Spec } R$  to a topological space by defining the closed sets of  $R$  to be the sets of the form  $V(I) = \{\mathfrak{p} : I \subseteq \mathfrak{p}\}$ . This defines a topology because

$$V(I) \cap V(J) = V(I+J) \quad \text{and} \quad V(I) \cup V(J) = V(I \cap J).$$

**Definition 1.13 (radical).** Let  $R$  be a ring. Given any  $I \subseteq R$ , set

$$\sqrt{I} = \{x \in R : x^n \in I \text{ for some } n \text{ depending on } x\}.$$

We call this the radical of  $I$ .

**Example 1.11.** We have  $\sqrt{(4)} = (2)$ .

**Example 1.12.** We have  $\mathfrak{N}_R = \sqrt{(0)}$ .

**Proposition 1.14.** Let  $I$  and  $J$  be ideals of a ring  $R$ . The following are fun to check:

- (i)  $\sqrt{\sqrt{I}} = \sqrt{I}$
- (ii)  $\sqrt{IJ} = \sqrt{I \cap J} = \sqrt{I} \cap \sqrt{J}$
- (iii)  $\sqrt{I} = R$  if and only if  $I = R$
- (iv)  $\sqrt{I+J} = \sqrt{\sqrt{I} + \sqrt{J}}$
- (v) If  $\mathfrak{p}$  is prime, then  $\sqrt{\mathfrak{p}^n} = \mathfrak{p}$
- (vi)  $\sqrt{I} = \bigcap_{\substack{I \subseteq \mathfrak{p} \\ \mathfrak{p} \text{ prime}}} \mathfrak{p}$

**Proposition 1.15.** Let  $I$  and  $J$  be ideals of a ring  $R$ . Then,

$$\sqrt{I} + \sqrt{J} = R \text{ implies } I + J = A.$$

*Proof.* We have  $\sqrt{I+J} = \sqrt{\sqrt{I} + \sqrt{J}} = \sqrt{R} = R$  so  $I + J = R$ . □

**Example 1.13** (Atiyah and Macdonald p. 12 Question 15). Let  $A$  be a ring and let  $X$  be the set of all prime ideals of  $A$ . For each subset  $E$  of  $A$ , let

$V(E)$  denote the set of all prime ideals of  $A$  which contain  $E$ .

Prove the following:

- (a) If  $\mathfrak{a}$  is the ideal generated by  $E$ , then  $V(E) = V(\mathfrak{a}) = V(r(\mathfrak{a}))$  (here,  $r(\mathfrak{a})$  denotes the radical of the ideal generated by  $\mathfrak{a}$  in the ring);
- (b)  $V(0) = X, V(1) = \emptyset$ ;
- (c) If  $(E_i)_{i \in I}$  is any family of subsets of  $A$ , then

$$V\left(\bigcup_{i \in I} E_i\right) = \bigcap_{i \in I} V(E_i);$$

- (d)  $V(\mathfrak{a} \cap \mathfrak{b}) = V(\mathfrak{a}\mathfrak{b}) = V(\mathfrak{a}) \cup V(\mathfrak{b})$  for any ideals  $\mathfrak{a}, \mathfrak{b}$  of  $A$ .

*Solution.*

- (a) By definition,  $V(E) = \{\mathfrak{p} \in X : E \subseteq \mathfrak{p}\}$ . Let  $\mathfrak{a}$  be the ideal generated by  $E$ . Then,  $\mathfrak{a}$  is the smallest ideal of  $A$  containing  $E$ . Hence,

$\mathfrak{p}$  contains  $E$  if and only if  $\mathfrak{p}$  contains  $\mathfrak{a}$ .

As such,  $V(E) = V(\mathfrak{a})$ . We then prove that  $V(\mathfrak{a}) = V(r(\mathfrak{a}))$ . Recall Definition 1.13 which states that  $r(\mathfrak{a}) = \{x \in A : x^n \in \mathfrak{a} \text{ for some } n\}$ . By (i) of Proposition 1.14,  $\mathfrak{p}$  is closed under taking radicals so a prime ideal  $\mathfrak{p}$  contains  $\mathfrak{a}$  if and only if it contains  $r(\mathfrak{a})$  and the result follows.

- (b) We have

$$V(0) = \{\mathfrak{p} \in A : 0 \in \mathfrak{p}\} \text{ and } V(1) = \{\mathfrak{p} \in A : 1 \in \mathfrak{p}\}.$$

So,  $V(0)$  contains all prime ideals  $\mathfrak{p}$  such that  $0 \in \mathfrak{p}$ . This is clearly  $X$ . Also, no prime ideal contains 1 as 1 generates the entire ring  $A$ . It follows that  $V(1) = \emptyset$ .

- (c) We first prove the forward inclusion. Let

$$\mathfrak{p} \in V\left(\bigcup_{i \in I} E_i\right) \text{ so } \bigcup_{i \in I} E_i \in \mathfrak{p}.$$

So,  $E_i \subseteq \mathfrak{p}$  for all  $i \in I$ , and it follows that  $\mathfrak{p}$  is contained in the intersection. The proof of the reverse inclusion is similar.

- (d) For any prime ideal  $\mathfrak{p}$ , it contains the ideal  $\mathfrak{a} \cap \mathfrak{b}$  if and only if it contains the ideals  $\mathfrak{a}$  and  $\mathfrak{b}$  of  $A$ , or equivalently  $\mathfrak{a}\mathfrak{b}$  by (i) of Proposition 1.10 since  $\mathfrak{a}\mathfrak{b} \subseteq \mathfrak{a} \cap \mathfrak{b}$  and both ideals generate the same radical in this case. So,  $V(\mathfrak{a} \cap \mathfrak{b}) = V(\mathfrak{a}\mathfrak{b})$ .

Next, we note that a prime ideal  $\mathfrak{p}$  contains  $\mathfrak{a}\mathfrak{b}$  if and only if  $\mathfrak{p}$  contains either  $\mathfrak{a}$  or  $\mathfrak{b}$ . This follows from the definition of a prime ideal. Hence,  $V(\mathfrak{a} \cap \mathfrak{b}) = V(\mathfrak{a}) \cup V(\mathfrak{b})$ .  $\square$

**Remark 1.3.** The results in Example 1.13 show that the sets  $V(E)$  satisfy the axioms for closed sets in a topological space. The resulting topology is called the *Zariski topology*. The topological space  $X$  is called the *prime spectrum of  $A$* , and is written  $\text{Spec}(A)$ .

**Example 1.14** (Atiyah and Macdonald p. 12 Question 17). For each  $f \in A$ , let  $X_f$  denote the complement of  $V(f)$  in  $X = \text{Spec}(A)$ . The sets  $X_f$  are open. Show that they form a basis of open sets for the Zariski topology, and that

- (i)  $X_f \cap X_g = X_{fg}$ ;
- (ii)  $X_f = \emptyset$  if and only if  $f$  is nilpotent;
- (iii)  $X_f = X$  if and only if  $f$  is a unit;
- (iv)  $X_f = X_g$  if and only if  $r((f)) = r((g))$  (here,  $r((f))$  denotes the radical of the ideal generated by  $f$  in the ring);
- (v)  $X$  is quasi-compact, i.e. every open covering of  $X$  has a finite subcovering;
- (vi) More generally, each  $X_f$  is quasi-compact
- (vii) An open subset of  $X$  is quasi-compact if and only if it is a finite union of sets  $X_f$ . The sets  $X_f$  are called *basic open sets* of  $X = \text{Spec}(A)$

*Hint:* To prove (v), remark that it is enough to consider a covering of  $X$  by basic open sets  $X_{f_i}$ , where  $i \in I$ . Show that the  $f_i$  generate the unit ideal and hence that there is an equation of the form

$$1 = \sum_{i \in J} g_i f_i \quad g_i \in A$$

where  $J$  is some finite subset of  $I$ . Then the  $X_{f_i}$ , where  $i \in J$ , cover  $X$ .

*Solution.* We first show that the collection of  $X_f$  forms a basis of open sets for the Zariski topology. Given a ring  $A$ , let  $f \in A$  and define

$$X_f = \{\mathfrak{p} \in \text{Spec}(A) : f \notin \mathfrak{p}\}.$$

For any ideal  $I \subseteq A$ , we define  $V(I) = \{\mathfrak{p} \in \text{Spec}(A) : I \subseteq \mathfrak{p}\}$  as the closed sets. The complement of  $V(f)$  is  $X_f$ , which as mentioned, is open. So, any open set in the Zariski topology is the union of such complements. Hence, the  $X_f$  form a basis.

- (i) By definition,

$$X_f \cap X_g = \{\mathfrak{p} \in \text{Spec}(A) : f \notin \mathfrak{p} \text{ and } g \notin \mathfrak{p}\} = \{\mathfrak{p} \in \text{Spec}(A) : fg \notin \mathfrak{p}\} = X_{fg}$$

and the result follows.

- (ii) For the forward direction, suppose  $X_f = \emptyset$ . Then,  $f \in \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Spec}(A)$ , so  $f$  is nilpotent. For the reverse direction, if  $f$  is nilpotent, there exists  $n \in \mathbb{N}$  such that  $f^n = 0$ . So, for every prime ideal  $\mathfrak{p}$ , we have  $f \in \mathfrak{p}$  so  $X_f = \emptyset$ .

- (iii) If  $f$  is a unit, then  $f \notin \mathfrak{p}$  for all  $\mathfrak{p} \in \text{Spec}(A)$ , so  $X_f = X$ . For the forward direction, if  $X_f = X$ , then  $f \notin \mathfrak{p}$  for all  $\mathfrak{p}$ . Hence,  $f$  is not contained in any maximal ideal, which implies  $f$  is a unit.
- (iv) Equivalent to saying that  $V(f) = V(g)$ .
- (v) Let  $\mathcal{S} = \{X_{f_i} : i \in I\}$  be an open cover of  $X$ . Since

$$X = \bigcup_{i \in I} X_{f_i} \quad \text{it implies} \quad \text{the } f_i \text{ generate the unit ideal } 1 = \sum_{i \in J} g_i f_i \text{ for some finite } J \subseteq I.$$

The result follows.

- (vi) Note that  $X_f$  can be covered by open sets  $X_{f_i}$  so we then apply the same argument as (v).
- (vii) Trivial. □

Recall from Definition 1.11 that a ring  $R$  is reduced if there exists no non-zero nilpotent elements, i.e.  $\mathfrak{N}_A = (0)$ . As such, we have the following proposition.

**Proposition 1.16.** For  $I \subseteq R$ ,

$$R/I \text{ is reduced} \quad \text{if and only if} \quad I = \sqrt{I}, \text{ i.e. } I \text{ is a radical ideal.}$$

**Definition 1.14 (Jacobson radical).** Given a ring  $R$ , define

$$J(R) = \bigcap_{\substack{\mathfrak{m} \subseteq R \\ \mathfrak{m} \text{ maximal}}} \mathfrak{m}.$$

In other words, the Jacobson radical of  $R$  is the intersection of all maximal ideals  $\mathfrak{m}$ .

**Proposition 1.17.**  $\mathfrak{N}_R \subseteq J(R)$

**Proposition 1.18.** We have

$$x \in J(R) \quad \text{if and only if} \quad 1 + yx \text{ is a unit for all } y \in R.$$

*Proof.* For the forward direction, choose  $x \in J(R)$ . Suppose on the contrary that  $1 + xy$  is not a unit. Then,  $1 + xy \in \mathfrak{m}$  for some maximal ideal  $\mathfrak{m}$ . As  $x \in \mathfrak{m}$ , then  $xy \in \mathfrak{m}$ , so  $1 \in \mathfrak{m}$ , which is a contradiction.

For the reverse direction, suppose on the contrary that  $x \notin \mathfrak{m}$  for some maximal ideal  $\mathfrak{m}$ . Then,  $(x) + \mathfrak{m} = R$ , so we can write  $1 = rx + m$  for some  $m \in \mathfrak{m}$ . Then,  $m = 1 - rx$  is not a unit. The result follows. □

**Definition 1.15 (ring of polynomials).** Given a ring  $R$ , consider the polynomial ring in one variable  $X$ , denoted by  $R[X]$ . It is defined as follows:

$$R[X] = \{ \text{polynomials } \sum r_i X^i : r_i \in R \text{ and } r_i = 0 \text{ for sufficiently large } i \}$$

Polynomial addition and multiplication (Cauchy product) are defined the obvious way.

**Definition 1.16 (formal Laurent series).** Let  $R$  be a ring. The ring of formal Laurent series in the variable  $X$  over  $R$  (often denoted by  $R[[X]]$ ) is defined as follows:

$$R[[X]] = \left\{ \sum_{i=N}^{\infty} r_i X^i : N \in \mathbb{Z}, r_i \in R \text{ for all } i, \text{ finitely many negative indices } i \text{ for which } r_i \neq 0 \right\}.$$

In other words, although the sum can extend infinitely in the positive direction, it can only extend finitely in the negative direction.

Definition 1.15 can be generalised to multiple indeterminates.

**Example 1.15 (construction of  $\mathbb{C}$  by taking quotient of maximal ideal).**  $\mathbb{R}[x]/(x^2 + 1) = \mathbb{C}$

**Example 1.16 (Gaussian integers).** Let

$$\mathbb{Z}[i] = \{a + bi : a, b \in \mathbb{Z}\} \quad \text{denote the set of Gaussian integers.}$$

Then,  $\mathbb{Z}[x]/(x^2 + 1) = \mathbb{Z}[i]$ .

**Example 1.17.** Consider  $(5) \subseteq \mathbb{Z}$ . Then

$$[i]/(5) = [x]/(x^2 + 1, 5) = \mathbb{F}_5[x]/(x^2 + 1) = \mathbb{F}_5[X]/((x-2)(x-3)) = \mathbb{F}_5 \times \mathbb{F}_5,$$

which is not an integral domain! Here,  $\mathbb{F}_5$  is the finite field of 5 elements. Therefore,  $(5)[i] \subseteq [i]$  is not prime<sup>†</sup>.

**Definition 1.17 (local ring).** A ring  $R$  is local if it has a unique maximal ideal  $\mathfrak{m}$ .

**Example 1.18.** Fields are local rings. To see why, the only ideals of any field  $F$  are  $\{0\}$  and  $F$ . Since  $\{0\}$  is the only proper ideal in  $F$ , it is the unique maximal ideal.

**Proposition 1.19.** If  $k$  is an arbitrary field, then

$$k[[X]] \text{ is a local ring as its only maximal ideal is } (X).$$

*Proof.* To show that any  $f \notin (X)$  is invertible, write  $f$  as

$$f = r_0 + Xg, \quad \text{where } r_0 \neq 0 \text{ and } g \in k[[X]]$$

We need to find  $h \in k[[X]]$  such that  $f \cdot h = 1$ . Using formal power series, define

$$h = \frac{1}{r_0 + Xg}.$$

Using the geometric series expansion, this can be rewritten as

$$h = \frac{1}{r_0} \cdot \frac{1}{1 + Xg/r_0} = \frac{1}{r_0} \sum_{i=0}^{\infty} \left( -\frac{Xg}{r_0} \right)^i.$$

Since  $Xg \in k[[X]]$ , the series converges in the formal sense, and we obtain

$$h = r_0^{-1} \sum_{i=0}^{\infty} X^i g^i r_0^{-i}.$$

Thus,  $h$  is a formal power series and  $f \cdot h = 1$ , proving that  $f$  is invertible. □

<sup>†</sup>Prof. David Hansen mentioned that he did not want to delve too deep into MA5202 with the introduction of number fields, etc.

**Example 1.19** (Atiyah and Macdonald p. 11 Question 10). Let  $A$  be a ring and  $\mathfrak{N}$  be its nilradical. Show that the following are equivalent:

- (i)  $A$  has exactly one prime ideal;
- (ii) every element of  $A$  is either a unit or nilpotent;
- (iii)  $A/\mathfrak{N}$  is a field

*Solution.* Recall from Proposition 1.6 that we defined the nilradical to be the set of nilpotent elements of  $A$ . We first prove that (i) implies (ii). Consider a maximal ideal of  $A$ , which must be prime, say  $\mathfrak{p}$ , since  $A$  has exactly one prime ideal. By Definition 1.17,  $A$  is a local ring. So, every element of  $A$  is a unit or nilpotent.

To prove (ii) implies (iii), it suffices to show that every element of  $A/\mathfrak{N}$  is invertible. Take any  $x + \mathfrak{N} \in A/\mathfrak{N}$  that is non-zero. So,  $x \notin \mathfrak{N}$ , i.e.  $x$  is not nilpotent. As such,  $x$  is a unit in  $A$ . Hence, there exists  $y \in A$  such that  $xy = 1$ . In  $A/\mathfrak{N}$ , this means that

$$(x + \mathfrak{N})(y + \mathfrak{N}) = xy + \mathfrak{N} = 1 + \mathfrak{N}.$$

Hence,  $x + \mathfrak{N}$  is invertible in  $A/\mathfrak{N}$ .

Lastly, we prove (iii) implies (i). Suppose  $A/\mathfrak{N}$  is a field. As such, the nilradical is maximal, and thus prime. As

$$\mathfrak{N} = \bigcap_{\mathfrak{p} \in \text{Spec} A} \mathfrak{p} \quad \text{it implies} \quad \text{every prime ideal contains } \mathfrak{N}.$$

Since  $\mathfrak{N}$  is maximal, then every prime ideal coincides with  $\mathfrak{N}$ . We conclude that  $A$  only has one prime ideal.  $\square$

**Example 1.20** (Atiyah and Macdonald p. 44 Question 5). Let  $A$  be a ring. Suppose that, for each prime ideal  $\mathfrak{p}$ , the local ring  $A_{\mathfrak{p}}$  has no nilpotent element  $\neq 0$ . Show that  $A$  has no nilpotent element  $\neq 0$ . If each  $A_{\mathfrak{p}}$  is an integral domain, is  $A$  necessarily an integral domain?

*Solution.* Suppose  $A$  has a non-zero nilpotent element  $x$ . Then,  $x$  belongs to all prime ideals  $\mathfrak{p}$  of  $A$ , and so do all of its powers  $x^n$ , for every  $n \in \mathbb{N}$ . Let  $\mathfrak{p}$  be a prime ideal. Then,  $(x/1) \in A_{\mathfrak{p}}$  is nilpotent. As such, for every  $\mathfrak{p}$ ,  $x \in \mathfrak{p}$  so  $x$  belongs to the intersection of all prime ideals of  $A$ . As such,  $\text{Spec}(A) = \mathfrak{N}_A$ . However, this contradicts the fact that  $A_{\mathfrak{p}}$  has no non-zero nilpotent elements.

The second part is false. Take  $A = \mathbb{Z}/6\mathbb{Z}$  which is not an integral domain. The prime ideals of  $A$  are  $\mathfrak{p}_2 = (2/6)$  and  $\mathfrak{p}_3 = (3/6)$  which correspond to 2 and 3 in  $\mathbb{Z}$ . We can then construct the local rings

$$A_{\mathfrak{p}_2} \cong \mathbb{Z}/2\mathbb{Z} \quad \text{and} \quad A_{\mathfrak{p}_3} \cong \mathbb{Z}/3\mathbb{Z} \quad \text{which are integral domains as they are fields.}$$

So, the second part is indeed false.  $\square$

## 1.2 Module Theory

**Definition 1.18** ( $R$ -module). Let  $R$  be a ring. An  $R$ -module  $M$  is an Abelian group  $(M, +, 0)$  equipped with a map of sets

$$R \times M \rightarrow M \quad \text{where} \quad (r, m) \mapsto m$$

such that the following properties hold:

- (i)  $(r_1 + r_2)m = r_1m + r_2m$

- (ii)  $r(r'm) = (rr')m$
- (iii)  $r(m_1 + m_2) = rm_1 + rm_2$
- (iv)  $1_R \cdot m = m$

**Definition 1.19** (*R-module homomorphism*). Given  $R$ -modules  $M$  and  $N$ , we have an obvious notion of an  $R$ -module homomorphism  $f : M \rightarrow N$ . Given any such  $f$ , we can generate some new  $R$ -modules, namely

$$\ker f \subseteq M \quad \text{im } f \subseteq N \quad \subseteq N \rightarrow \text{coker } f.$$

**Example 1.21.** An ideal  $I \subseteq R$  is an  $R$ -submodule of  $R$ .

**Example 1.22.** Let  $M$  and  $N$  be  $R$ -modules. Then,

$$\text{Hom}_R(M, N) = \{R\text{-module maps } f : M \rightarrow N\}.$$

This is a natural  $R$ -module as

$$(f_1 + f_2)(m) = f_1(m) + f_2(m) \quad \text{and} \quad (rf)(m) = f(rm) = rf(m).$$

**Example 1.23.** We have  $\text{Hom}_R(R, M) = M$  by sending  $f \mapsto f(1)$  and  $f(1) \mapsto (r \mapsto rm)$ .

**Example 1.24.** Given  $I \subseteq R$ , we have  $\text{Hom}_R(R/I, M) = M[I]$ . Here,  $M[I]$  refers to the torsion submodule of  $M$  associated with  $I$ , where we define

$$M[I] = \{m \in M : \text{there exists } i \in I \text{ such that } im = 0\}.$$

**Definition 1.20** (*submodule*). For a ring  $R$  with an ideal  $I \subseteq R$ , and an  $R$ -module  $M$ ,  $IM$  denotes the submodule of  $M$  generated by the expressions of the form  $i_1m_1 + \cdots + i_jm_j$ .

**Example 1.25.** If  $M = R$  then we have  $IR = I$ .

# Chapter 2

## Basic Commutative Algebra

### 2.1

#### Exact Sequences of Modules

**Definition 2.1** (complex and exact sequences). Fix a ring  $R$ . A sequence of  $R$ -module homomorphisms

$$\dots \xrightarrow{f_{i-1}} M_i \xrightarrow{f_i} M_{i+1} \xrightarrow{f_{i+1}} M_{i+2} \xrightarrow{f_{i+2}} \dots$$

- (i) is *complex* if  $\text{im } f_i \subseteq \ker f_{i+1}$  for all  $i$ , i.e.  $f_{i+1} \circ f_i = 0$  for all  $i$ ;
- (ii) is an *exact sequence* if  $\text{im } f_i = \ker f_{i+1}$

We will often be in a situation where  $M_i = 0$  for all but finitely many  $i$ .

**Example 2.1.** In the sequence of  $R$ -module homomorphisms  $0 \rightarrow M \xrightarrow{f} N$ ,  $f$  is injective as  $\ker f = \{0\}$ .

**Example 2.2.** In the sequence of  $R$ -module homomorphisms  $N \xrightarrow{g} Q \rightarrow 0$ ,  $g$  is surjective.

**Example 2.3.** The sequence of  $R$ -module homomorphisms

$$0 \rightarrow M \xrightarrow{\text{id}} 0 \quad \text{is always exact.}$$

**Definition 2.2** (short exact sequence). Suppose the sequence of  $R$ -module homomorphisms

$$0 \rightarrow M \xrightarrow{f} N \xrightarrow{g} Q \rightarrow 0 \quad \text{is exact.}$$

This is equivalent to saying that  $f$  is injective,  $g$  is surjective, and  $\text{im } f = \ker g$ .

**Example 2.4.** Consider the following sequence of Abelian groups:

$$\mathbb{Z} \xrightarrow{\times 2} \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$$

The first homomorphism maps each element  $i \in \mathbb{Z}$  to the element  $2i \in \mathbb{Z}$ . The second homomorphism maps each element  $i$  in  $\mathbb{Z}$  to the quotient group  $\mathbb{Z}/2\mathbb{Z}$ , that is  $j \equiv i \pmod{2}$ . This is an exact sequence since the image of the **red homomorphism** is the kernel of the **blue homomorphism**<sup>†</sup>.

**Proposition 2.1.** Given any  $R$ -module homomorphism  $f : M \rightarrow N$ , we can always obtain the following two short exact sequences:

$$\begin{aligned} 0 \rightarrow \ker f \rightarrow M \rightarrow \text{im } f \rightarrow 0 \\ 0 \rightarrow \text{im } f \rightarrow N \rightarrow \text{coker } f \rightarrow 0 \end{aligned}$$

Recall that  $\text{coker } f$  measures how far  $f$  is from being surjective. It is defined as the quotient module  $f/\text{im } f$ .

<sup>†</sup>In Category Theory *language*, the hook arrow  $\hookrightarrow$  denotes an injective homomorphism so we say that it is a monomorphism; the two-headed arrow  $\twoheadrightarrow$  is a surjective homomorphism so we say that it is an epimorphism.



**Definition 2.3** (finitely generated module). An  $R$ -module  $M$  is finitely generated if there exist elements  $x_1, \dots, x_n \in M$  such that all  $m \in M$  can be expressed as a finite linear combination, i.e.

$$\sum_{i=1}^n r_i m_i \quad \text{for some } r_i \in R.$$

Note that if  $M$  is a finitely generated  $R$ -module, it is equivalent to saying that there exists an exact sequence

$$\begin{aligned} R^n &\rightarrow M \rightarrow 0 \\ e_i &\mapsto x_i \end{aligned}$$

**Definition 2.4** (finitely presented module). An  $R$ -module  $M$  is finitely presented if there exists an exact sequence

$$R^m \rightarrow R^n \rightarrow M \rightarrow 0 \quad \text{for some } m, n \in \mathbb{N}.$$

**Example 2.5.** Let  $k$  be a field. Define

$$R = k[x_1, x_2, x_3, \dots] \quad \text{and} \quad \mathfrak{m} = (x_1, x_2, x_3, \dots) \quad \text{so } M = R/\mathfrak{m} \cong k.$$

Then,  $M$  is finitely generated but not finitely presented as  $\mathfrak{m}$  is not finitely generated as an  $R$ -module.

**Example 2.6.** Suppose we have a short exact sequence  $0 \rightarrow M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$ . Then,  $M_1$  and  $M_3$  are finitely generated, which implies  $M_2$  is also finitely generated (in Example 2.7, we will discuss the proof of this result but for the case where the sequence is exact, i.e. no assumption of it being a short exact sequence). The same result holds if we change the term ‘finitely generated’ to ‘finitely presented’.

**Example 2.7** (Atiyah and Macdonald p. 32 Question 9). Let

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0 \quad \text{be an exact sequence of } A\text{-modules.}$$

If  $M'$  and  $M''$  are finitely generated, then so is  $M$ .

*Solution.* Suppose

$$M' \text{ is generated by } x_1, \dots, x_n \quad \text{and} \quad M'' \text{ is generated by } z_1, \dots, z_m.$$

Suppose  $u : M' \rightarrow M$  and  $v : M \rightarrow M''$  are  $A$ -module homomorphisms. Let  $v(y_i) = z_i$  for all  $1 \leq i \leq m$ . Also, let  $x \in M$ . Then, there exist  $b_1, \dots, b_m \in A$  such that  $v(x) = b_1 z_1 + \dots + b_m z_m$ . Hence,

$$v(x) = b_1 v(y_1) + b_m v(y_m) \quad \text{so} \quad v(x) = v(b_1 y_1 + \dots + b_m y_m).$$

Hence,  $x - b_1 y_1 - \dots - b_m y_m \in \ker v$ . As the sequence is exact, then  $\text{im } u = \ker v$ . So, there exist  $a_1, \dots, a_n \in A$  such that

$$\begin{aligned} x - (b_1 y_1 + \dots + b_m y_m) &= a_1 u(x_1) + a_n u(x_n) \\ x &= b_1 y_1 + \dots + b_m y_m + a_1 u(x_1) + a_n u(x_n) \end{aligned}$$

so  $M$  is generated by  $u(x_1), \dots, u(x_n), y_1, \dots, y_m$ . □

**Example 2.8** (Atiyah and Macdonald p. 32 Question 12). Let  $M$  be a finitely generated  $A$ -module and  $\varphi : M \rightarrow A^n$  be a surjective homomorphism. Show that  $\ker \varphi$  is finitely generated.

*Hint:* Let  $e_1, \dots, e_n$  be a basis for  $A^n$  and choose  $u_i \in M$  such that  $\varphi(u_i) = e_i$  for all  $1 \leq i \leq n$ . Show that  $M$  is the direct sum of  $\ker \varphi$  and the submodule generated by  $u_1, \dots, u_n$ .

*Solution.* Let  $m \in M$ , so  $\varphi(m) \in A^n$ . We can write

$$\varphi(m) = a_1 e_1 + \dots + a_n e_n \quad \text{where } a_1, \dots, a_n \in A.$$

Also, let  $U$  be a submodule of  $M$ . Since  $M$  is finitely generated, then  $U$  is also finitely generated by say  $u_1, \dots, u_n$ . So, there exist  $a_1, \dots, a_n \in A$  such that

$$\begin{aligned} u &= a_1 u_1 + \dots + a_n u_n \\ \varphi(u) &= a_1 \varphi(u_1) + \dots + a_n \varphi(u_n) \\ &= a_1 e_1 + \dots + a_n e_n \end{aligned}$$

Since the RHS is  $\varphi(m)$ , then  $\varphi(u - m) = 0$ , so  $u - m \in \ker \varphi$ . Thus, for any  $m \in M$ , we can decompose it as  $m = (m - u) + u$ , which shows that  $M$  is the sum of  $\ker \varphi$  (elements of the form  $m - u$ ) and the submodule generated by  $u_1, \dots, u_n$ .

We then show that the sum is direct, i.e.  $\ker \varphi \cap U = \emptyset$ . Suppose  $m \in \ker \varphi \cap U$ . Then,  $m \in \ker \varphi$  and  $m \in U$ . The former tells us that  $\varphi(m) = 0$ , whereas the latter tells us that

$$m = a_1 u_1 + \dots + a_n u_n \quad \text{where } a_1, \dots, a_n \in A.$$

Applying  $\varphi$  to both sides, we obtain  $0 = a_1 e_1 + \dots + a_n e_n$ . Since  $e_1, \dots, e_n$  is a basis for  $A^n$ , then  $a_1 = \dots = a_n = 0$ . Hence  $m = 0$  and the result follows. □

**Lemma 2.1** (snake lemma). Suppose we are given a commutative diagram of  $R$ -modules

$$\begin{array}{ccccccccc} 0 & \longrightarrow & M' & \xrightarrow{a} & M & \xrightarrow{b} & M'' & \longrightarrow & 0 \\ & & \downarrow f' & & \downarrow f & & \downarrow f'' & & \\ 0 & \longrightarrow & N' & \xrightarrow{c} & N & \xrightarrow{d} & N'' & \longrightarrow & 0 \end{array}$$

where the rows are exact. Then, there exists a natural exact sequence

$$0 \rightarrow \ker f' \rightarrow \ker f \rightarrow \ker f'' \xrightarrow{\delta} \operatorname{coker} f' \rightarrow \operatorname{coker} f \rightarrow \operatorname{coker} f'' \rightarrow 0.$$

*Proof.* It suffices to construct  $\delta$ . Given  $x \in M''$  such that  $f''(x) = 0$ , pick  $y \in M$  such that  $b(y) = x$ . Then  $0 = f''(x) = f''(b(y)) = d(f(y))$ . Thus  $f(y) \in \ker d = \operatorname{im} c$ , so there exists a unique  $z \in N'$  such that  $c(z) = f(y)$ . We set  $\delta(x) = z + f'$ .

For well-definedness, we need to see that the choice of  $y$  does not matter. If  $y' \in M$  with  $b(y') = x$ , then  $y - y' \in \ker b = \operatorname{im} a$  so  $f(y) - f(y') \in \operatorname{im} c$ . □

**Proposition 2.2** (Nakayama's lemma, useless version). Fix a ring  $A$ . Pick  $M$  to be a finitely generated  $A$ -module and  $I \subseteq A$  be an ideal. Let  $\varphi : M \rightarrow M$  be an  $A$ -module homomorphism such that  $\varphi(M) \subseteq IM$ .

Then,

there exists an equation  $\varphi^n + a_1\varphi^{n-1} + a_2\varphi^{n-2} + \dots + a_n = 0$  where  $a_i \in I$ .

*Proof.* Pick  $x_1, \dots, x_n \in M$  generating  $M$ . Then,  $\varphi(x_i) \in IM$ . Since  $M$  is finitely generated, then

$$\varphi(x_i) = \sum_{j=1}^n a_{ij}x_j \quad \text{for some choice of } a_{ij} \in I.$$

We can write the equation as

$$\sum_{j=1}^n (\delta_{ij}\varphi(x_i) - a_{ij})x_j = 0.$$

Write the above as  $\mathbf{A}\mathbf{x} = \mathbf{0}$  so

$$A_{ij} = \delta_{ij}\varphi(x_i) - a_{ij} \quad \text{and} \quad \mathbf{x} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}.$$

Recall from MA2001 that

$$\det(\mathbf{A})\mathbf{I}_n = \text{adj}(\mathbf{A})\mathbf{A} \quad \text{where} \quad \text{adj}(\mathbf{A})_{ij} = (-1)^{i+j}M_{ji}.$$

Hence,

$$\begin{bmatrix} \det(\mathbf{A})x_1 \\ \vdots \\ \det(\mathbf{A})x_n \end{bmatrix} = \det(\mathbf{A})\mathbf{I}_n\mathbf{x} = \text{adj}(\mathbf{A})\mathbf{A}\mathbf{x} = \mathbf{0}.$$

As such,  $\det(\mathbf{A})x_i = 0$  for all  $1 \leq i \leq n$ . So,  $\det(\mathbf{A}) = 0$  in  $\text{Hom}_A(M, M)$ . We conclude that  $\det(\mathbf{A}) = \varphi^n + a_1\varphi^{n-1} + \dots + a_n$ , where  $a_i \in I$ . □

**Corollary 2.1.** Let  $M$  be a finitely generated  $A$ -module and  $I \subseteq A$  be an ideal such that  $IM = M$ . Then,

there exists  $a \equiv 1 \pmod{I}$  such that  $aM = 0$ .

*Proof.* If  $IM = M$ , then by Proposition 2.2, we have  $0 = 1 + a_1 + \dots + a_n$  as elements of  $\text{Hom}_A(M, M)$ , where the RHS is an element of  $I$ . Setting  $a = 1 + a_1 + \dots + a_n$ , the result follows. □

**Proposition 2.3 (Nakayama's lemma V1).** Let  $M$  be a finitely generated  $A$ -module and  $I \subseteq J(A)$  be an ideal (recall that  $J(A)$  is the Jacobson radical of  $A$ ). If  $IM = M$ , then  $M = 0$ .

*Proof.* By Corollary 2.1, we obtain some  $a = 1 + I$  with  $aM = 0$ . However,  $I \subseteq J(A)$ , which implies  $a \in A^*$ . So,  $M = a^{-1}(aM) = 0$ . □

**Example 2.9.** Let  $A = \mathbb{Z}/4\mathbb{Z}$  and  $M$  be a finitely-generated  $A$ -module. Recall that the Jacobson radical  $J(A)$  is the intersection of all maximal ideals  $\mathfrak{m}$  of  $A$ , for which there is only one  $(2)$ . As such,  $J(A) = 2A$ . Setting  $I = J(A)$ , we have  $J(A)M = M$  since  $2M = M$ . As such,

$$IM = M \quad \text{which implies} \quad M = 0 \text{ (the zero module).}$$

**Example 2.10** (Atiyah and Macdonald p. 32 Question 10). Let  $A$  be a ring,  $\mathfrak{a}$  an ideal contained in  $J(A)$ ; let  $M$  be an  $A$ -module and  $N$  a finitely generated  $A$ -module, and let  $u : M \rightarrow N$  be a homomorphism. If  $M/\mathfrak{a}M \rightarrow N/\mathfrak{a}N$  is surjective, prove that  $u$  is surjective.

*Solution.* We will make use of V1 of Nakayama’s lemma (Proposition 2.3). Define  $L = N/u(M)$ . We shall prove that  $L = 0$ , i.e.  $N = u(M)$ , and consequently,  $u$  is surjective. Since

$$M/\mathfrak{a}M \rightarrow N/\mathfrak{a}N \text{ is surjective,}$$

then for every element  $\bar{n} \in N/\mathfrak{a}N$ , there exists  $\bar{m} \in M/\mathfrak{a}M$  mapping to it. As such,  $N/\mathfrak{a}N = (u(M) + \mathfrak{a}N)/\mathfrak{a}N$ , or equivalently,  $N = u(M) + \mathfrak{a}N$ . As such, we have

$$L = N/u(M) = (u(M) + \mathfrak{a}N)/u(M).$$

From here, one can deduce that  $L \subseteq \mathfrak{a}N/u(M)$ , so  $L = \mathfrak{a}L$ . Since  $\mathfrak{a}$  is an ideal contained in the Jacobson radical  $J(A)$ , then by applying Nakayama’s lemma (Proposition 2.3) to the finitely-generated  $A$ -module  $L$  (since  $L = \mathfrak{a}L$ ), then  $L = 0$ . The result follows.  $\square$

**Proposition 2.4** (Nakayama’s lemma V2). Let  $M$  be a finitely generated  $A$ -module,  $N \subseteq M$  and  $I \subseteq J(A)$ . Then,

$$M = IM + N \text{ implies } M = N.$$

*Proof.* Applying version 1 of Nakayama’s lemma (Proposition 2.3) to  $Q = M/N$ , we obtain

$$IQ = (IM + N)/N = M/N = Q.$$

Again by applying Proposition 2.3, we have  $Q = 0$  so  $M = N$ .  $\square$

**Proposition 2.5** (Nakayama’s lemma V3). Let  $(A, \mathfrak{m})$  be a local ring and  $k = A/\mathfrak{m}$  denote its residue field. If  $M$  is a finitely generated  $A$ -module and  $x_1, \dots, x_n \in M/\mathfrak{m}M$  span  $M/\mathfrak{m}M$  as a  $k$ -vector space, then any choice of lifts  $\tilde{x}_1, \dots, \tilde{x}_n \in M$  generate  $M$  as an  $A$ -module.

*Proof.* Take  $N \subseteq M$  to be the submodule generated by  $\tilde{x}_1, \dots, \tilde{x}_n$ . Then,  $M = N + \mathfrak{m}M$ , so  $M = N$  by Proposition 2.5.  $\square$

**Example 2.11.** Recall that every field is a local ring (Example 1.18). For any field  $k$ , let  $A = k[x]$  and let  $\mathfrak{m} = (x)$  be the maximal ideal in  $A$ . Take  $M = A/(x^2)$  as an  $A$ -module. The residue field is  $k = A/\mathfrak{m}$ . The module  $M/\mathfrak{m}M = (A/(x^2))/(x) = k$ , which is a 1-dimensional  $k$ -vector space. Also, the element  $\bar{1} \in M/\mathfrak{m}M$  spans  $M/\mathfrak{m}M$  as a  $k$ -vector space. By Proposition 2.5, the lift  $\tilde{1} = 1 \in M$  generates  $M$  as an  $A$ -module. We conclude that  $A/(x^2)$  is cyclic as an  $A$ -module.

## 2.2 Localization

**Definition 2.5** (multiplicatively closed set). Fix a ring  $A$ . A subset  $S \subseteq A$  is said to be multiplicatively closed if

$$1 \in S \text{ and for all } s_1, s_2 \in S \text{ we have } s_1s_2 \in S.$$

**Example 2.12.** For any ring  $A$ , the set of non-zero divisors is multiplicatively closed.

**Example 2.13.** For any  $f \in A$ ,  $\{1, f, f^2, \dots\}$  is multiplicatively closed.

**Example 2.14.** For any prime ideal  $\mathfrak{p}$  of  $A$ , the set  $A \setminus \mathfrak{p}$  is multiplicatively closed.

**Theorem 2.1.** Given a ring  $A$  and any multiplicatively closed  $S \subseteq A$ , there exists a naturally associated ring  $S^{-1}A$  equipped with a ring homomorphism  $\varphi : A \rightarrow S^{-1}A$  ( $S^{-1}A$  denotes the localization of  $A$  at  $S$ ) such that for any ring homomorphism  $f : A \rightarrow B$  where  $f(S) \subseteq B^\times$ , there exists a *unique* ring homomorphism

$$f' : S^{-1}A \rightarrow B \quad \text{such that} \quad f = f' \circ \varphi.$$

Hence, the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow \varphi & \uparrow \exists! f' \\ & & S^{-1}A \end{array}$$

In other words words,  $\varphi_S : A \rightarrow S^{-1}A$  is *universal* for ring homomorphisms  $f : A \rightarrow B$  sending  $S$  to units.

*Proof.* We will first construct  $S^{-1}A$  as a set. Let

$$S^{-1}A = (A \times S) / \sim,$$

with  $(a, s) \sim (a', s')$  if and only if there exists  $t \in S$  such that  $t(as' - a's) = 0$ . We define

$$\begin{aligned} (a, s) \cdot (a', s') &= (aa', ss') \\ (a, s) + (a', s') &= (as' + a's, ss') \end{aligned}$$

The multiplicative identity is  $(1, 1)$  and the additive identity is  $(0, 1)$ .

We will write  $\frac{a}{s}$  for the equivalence class of  $(a, s)$ . The universal map  $\varphi_S : A \rightarrow S^{-1}A$  is defined by  $a \mapsto (a, 1)$ . Given  $f : A \rightarrow B$ , suppose that  $f = f' \circ \varphi_S$  for some  $f' : A \rightarrow S^{-1}A$ , then  $f(S) \subseteq f'((S^{-1}A)^\times) \subseteq B^\times$ .

Now suppose that  $f(S) \subseteq B^\times$ . Note that  $a \in \ker \varphi_S$  if and only if there exists  $s \in S$  such that  $sa = 1$ . Now define  $f'(\frac{a}{s}) = f(a)f(s)^{-1}$ . We need to show that this is well-defined, i.e. independent of the choice of representatives. If  $(a, s) \sim (a', s')$  then there exists  $t \in S$  such that  $tas' - ta's = 0$ , so applying  $f$  gives

$$f(t)f(a)f(s') = f(t)f(a')f(s) = 0,$$

whence multiplying by  $(f(s)f(s')f(t))^{-1}$  gives  $f(a)f(s)^{-1} - f(a')f(s')^{-1} = 0$  as required. It is clear by construction that  $f' \circ \varphi_S = f$ . This map is unique because  $\ker \varphi_S \subseteq \ker f$ . Note that  $\varphi_S(s)$  is a unit for all  $s \in S$ , since  $(s, 1) = (1, s) = (1, 1) = 1_{S^{-1}A}$ . □

**Corollary 2.2.** We have

$$\varphi_S : A \rightarrow S^{-1}A \text{ is an isomorphism if and only if } S \subseteq A^\times.$$

*Proof.* For the forward direction, note that  $\varphi(S) \subseteq (S^{-1}A)^\times$ , but  $\varphi_S$  is an isomorphism, so  $S \subseteq A^\times$ . For the reverse direction, we use the universal property of  $S^{-1}A$  on  $\text{id} : A \rightarrow A$  to find  $f^{-1} : S^{-1}A \rightarrow A$  such that  $\text{id} = f \circ \varphi_S$ . The result follows. □

We briefly remark that  $\varphi_S$  is not always injective. For instance, if  $A = \mathbb{Z}/6\mathbb{Z}$  and  $S = \{1, 2, 4\}$ , then  $S^{-1}A = \mathbb{Z}/3\mathbb{Z}$ . One checks that  $S \subseteq A$ . Moreover,  $S$  is a multiplicatively closed subset of  $A$ . That is to say,  $S$  is closed under multiplication. We will justify that  $S^{-1}A = \mathbb{Z}/3\mathbb{Z}$  (recall that this process is known as localization, which makes the elements of  $S$  invertible). Elements of  $S^{-1}A$  are of the form  $\frac{a}{s}$ , where  $a \in A$  and  $s \in S$ , with the rule that

$$\frac{a}{s} = \frac{b}{t} \text{ if and only if there exists a unit } u \text{ such that } u(sa - tb) = 0 \text{ in } A.$$

Consider  $2 \in S$ , which satisfies  $\gcd(2, 6) = 2$ . So, multiplication by 2 annihilates  $\bar{3}$ , i.e.  $2 \cdot \bar{3} = \bar{0}$ . The condition 2 is invertible in the localization implies that 3 must be sent to 0. As such, the ring  $\mathbb{Z}/6\mathbb{Z}$  effectively collapses as if we were also factoring the ideal generated by 3. Indeed, it is clear that 2 is invertible in  $\mathbb{Z}/3\mathbb{Z}$  since  $2 \cdot 2 \equiv 1 \pmod{3}$ .

Having said all the above, if however  $S$  does not contain any zero divisors, then  $\varphi_S$  is injective. In particular, if  $A$  is an integral domain, then  $\varphi_S$  is injective for any  $S$  and  $S^{-1}A$  is also an integral domain.

**Proposition 2.6.** If  $S \subseteq T \subseteq A$  are multiplicatively closed, then the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\varphi_S} & S^{-1}A \\ & \searrow \varphi_T & \downarrow \\ & & T^{-1}A \end{array} \quad \begin{array}{ccc} & & \varphi_{\varphi_S(T)} \\ & & \searrow \\ & & (\varphi_S(T))^{-1}S^{-1}A \\ & \dashrightarrow & \\ & & \end{array}$$

**Example 2.15.** Choose  $f \in A$  and take  $S = \{1, f, f^2, \dots\}$  which is multiplicatively closed. Then, we can write  $A_f = S^{-1}A$ .

**Proposition 2.7.** We have  $A_f \cong A[X]/(1 - fX)$ .

Now let  $R$  be a ring,  $S \subseteq R$  be multiplicatively closed, and consider  $\varphi_S : R \rightarrow S^{-1}R$ . If  $I \subseteq R$  is an ideal of  $R$ , then  $\varphi_S(I)S^{-1}R \subseteq S^{-1}R$  is an ideal of  $S^{-1}R$ . Likewise if  $J \subseteq S^{-1}R$  is an ideal of  $S^{-1}R$ , then  $\varphi_S^{-1}(J) \subseteq R$  is an ideal of  $R$ . We can verify the following facts:

- $\varphi_S^{-1}(\varphi_S(I)S^{-1}R) \supseteq I$ ;
- $J \supseteq \varphi_S(\varphi_S^{-1}(J))S^{-1}R$

In general, equality does not hold. But things are nicer with prime ideals.

**Theorem 2.2.** There exists a canonical bijection

$$\{\text{prime } \mathfrak{p} \subseteq R \mid \mathfrak{p} \cap S = \emptyset\} \cong \{\text{prime ideals } \mathfrak{q} \subseteq S^{-1}R\}$$

sending  $\mathfrak{p} \rightarrow \varphi_S(\mathfrak{p})S^{-1}R$  and  $\mathfrak{q} \mapsto \varphi_S^{-1}(\mathfrak{q})$ .

**Definition 2.6 (saturation).** Let  $\mathfrak{a} \subseteq R$  be any subset. We define the *saturation* of  $\mathfrak{a}$  with respect to  $S$  to be

$$\mathfrak{a}^S = \{a \in R \mid sa \in \mathfrak{a} \text{ for some } s \in S\}.$$

If  $\mathfrak{a} = \mathfrak{a}^S$ , we say that  $\mathfrak{a}$  is *saturated*.

**Proposition 2.8.** Let  $R$  be a ring. Fix a multiplicatively closed subset  $S \subseteq R$ . Then, the following hold:

- (i) If  $\mathfrak{b} \subseteq S^{-1}R$  is an ideal, then  $\varphi_S^{-1}(\mathfrak{b}) = (\varphi_S^{-1}(\mathfrak{b}))^S$  and  $\mathfrak{b} = \varphi_S^{-1}(\mathfrak{b})S^{-1}R$
- (ii) If  $\mathfrak{b} \subseteq R$  is an ideal, then  $\varphi_S(\mathfrak{a})S^{-1}R = \varphi_S(\mathfrak{a}^S)S^{-1}R$  and  $\varphi_S^{-1}(\varphi_S(\mathfrak{a})S^{-1}R) = \mathfrak{a}^S$
- (iii) Let  $\mathfrak{p} \subseteq R$  be a prime ideal with  $\mathfrak{p} \cap S = \emptyset$ . Then  $\mathfrak{p} = \mathfrak{p}^S$  and  $\varphi_S(\mathfrak{p})S^{-1}R \subseteq S^{-1}R$  is prime.

*Proof.* We first prove the first part of (i). Suppose  $a \in \varphi_S^{-1}(\mathfrak{b})$ . Then,

$$\frac{as}{1} \in \mathfrak{b} \subseteq S^{-1}R.$$

Since  $s$  is a unit in  $S^{-1}R$ , then we can write

$$\frac{a}{1} = \frac{as}{1} \cdot \frac{1}{s} \in \mathfrak{b} \quad \text{so} \quad \varphi_S(a) \in \mathfrak{b}.$$

As such,  $a \in \varphi_S^{-1}(\mathfrak{b})$ . One can deduce  $\subseteq$  of the first part from here.  $\supseteq$  is obvious.

We then prove the second part of (i). Suppose  $\varphi_S(a) \in \mathfrak{b}$ , so  $a \in \varphi_S^{-1}(\mathfrak{b})$ . So,

$$\frac{a}{s} = \frac{a}{1} \cdot \frac{1}{s} \in \varphi_S^{-1}(\mathfrak{b})S^{-1}R,$$

which implies  $\mathfrak{b} \subseteq \varphi_S^{-1}(\mathfrak{b})S^{-1}R$ , proving  $\subseteq$ . Note that  $\supseteq$  is obvious, so (i) holds.

We then prove the first part of (ii). Suppose  $a \in \mathfrak{a}^S$ , i.e. there exists  $s$  with  $sa \in \mathfrak{a}$ . Thus,

$$\frac{a}{1} = \frac{as}{1} \cdot \frac{1}{s} \in \varphi_S(\mathfrak{a})S^{-1}R.$$

Thus,  $\subseteq$  follows. Note that  $\supseteq$  is obvious, so the first part of (ii) follows. For the second part, suppose  $x \in \varphi_S^{-1}(\varphi_S(\mathfrak{a})S^{-1}R)$ . Then,

$$\frac{x}{1} = \frac{a}{s} \quad \text{with} \quad a \in \mathfrak{a} \text{ and } s \in S.$$

This implies that there exists  $t \in S$  such that  $xst = at$  in  $\mathfrak{a}$ . As such,  $x \in \mathfrak{a}^S$ . Thus,  $\varphi_S^{-1}(\varphi_S(\mathfrak{a})S^{-1}R) \subseteq \mathfrak{a}^S$ , proving the forward direction  $\subseteq$ . The reverse direction  $\supseteq$  holds as the left side is saturated by 1. As such, the second part of (ii) holds, so (ii) holds.

Lastly, we prove (iii). For the first part, take  $as \in \mathfrak{p}$  for some prime ideal  $\mathfrak{p} \subseteq R$ . Since  $\mathfrak{p} \cap S = \emptyset$ , this implies  $a \in \mathfrak{p}$ . As such,  $\mathfrak{p}^S \subseteq \mathfrak{p}$ , proving  $\supseteq$ . The proof of the forward direction  $\subseteq$  is clear.

We then prove the second part. Note that

$$\varphi_S(\mathfrak{p})S^{-1}R \neq S^{-1}R \quad \text{because} \quad \varphi_S^{-1}(\varphi_S(\mathfrak{p})S^{-1}R) = \mathfrak{p}^S = \mathfrak{p}.$$

The last equality follows from the first part of (iii). Now, suppose we are given some element

$$\frac{a}{s} \cdot \frac{b}{t} \in \varphi_S(\mathfrak{p})S^{-1}R.$$

Then,

$$ab \in \varphi_S^{-1}(\varphi_S(\mathfrak{p})S^{-1}R) = \mathfrak{p}.$$

Since  $\mathfrak{p}$  is a prime ideal, then either  $a \in \mathfrak{p}$  or  $b \in \mathfrak{p}$ . So,

$$\frac{a}{s} \text{ or } \frac{b}{t} \text{ is an element of } \varphi_S(\mathfrak{p})S^{-1}R.$$

It follows that  $\varphi_S(\mathfrak{p})S^{-1}R$  is prime, completing the proof.  $\square$

**Example 2.16.** If  $S = \{1, f, f^2, \dots\}$  with  $S^{-1}R = R_f$ , then the induced map  $\text{Spec}(R_f) \rightarrow \text{Spec}(R)$  is injective with image  $\{\mathfrak{p} \subseteq R \mid f \notin \mathfrak{p}\} = \text{Spec}(R \setminus V(f))$ . In particular, the image is an open subset.

**Definition 2.7 (localization).** Let  $A$  be a ring and  $S \subseteq A$  be multiplicatively closed. Suppose  $M$  is an  $A$ -module. Define an  $S^{-1}A$  module as follows:

$$S^{-1}M = M \times S / \sim \quad \text{where} \quad (m, s) \sim (m', s') \text{ if there exists } t \in S \text{ such that } t(s'm - sm') = 0$$

We use the notation  $\frac{m}{s}$  to refer to the equivalence class  $(m, s)$ . Addition and scalar multiplication are defined in the following obvious way:

$$(m, s) + (m', s') = (s'm + sm', ss') \quad \text{and} \quad (a, s) \cdot (m, t) = (am, st)$$

If  $f : M \rightarrow N$  is any map of  $A$ -modules, we obtain an induced map

$$S^{-1}f : S^{-1}M \rightarrow S^{-1}N \quad \text{of } S^{-1}A\text{-modules.}$$

In Category theory, we say that  $S^{-1}(\cdot)$  is a functor from an  $A$ -module to an  $S^{-1}A$ -module.

**Proposition 2.9.** If

$$M' \xrightarrow{f} M \xrightarrow{g} M'' \quad \text{is exact at } M,$$

then

$$S^{-1}M' \xrightarrow{S^{-1}f} S^{-1}M \xrightarrow{S^{-1}g} S^{-1}M'' \quad \text{is exact at } S^{-1}M.$$

*Proof.* Note that  $\text{im}(S^{-1}f) \subseteq \text{ker}(S^{-1}g)$  since

$$(S^{-1}g \circ S^{-1}f) \left( \frac{m'}{s} \right) = \frac{g(f(m'))}{s} = \frac{0}{s} = 0.$$

Next, suppose we are given some  $\frac{m}{s} \in S^{-1}M$  with

$$\frac{g(m)}{s} = 0 \quad \text{in } S^{-1}M''.$$

Then, there exists  $t \in S$  such that  $t(g(m)) = 0$  in  $M''$ . This implies  $g(tm) = 0$ . As such, there exists  $n \in M'$  such that  $f(n) = tm$ . Applying  $S^{-1}f$  yields

$$S^{-1} \left( \frac{n}{ts} \right) = \frac{f(n)}{ts} = \frac{tm}{ts} = \frac{m}{s}$$

which is contained in  $S^{-1}f$ . □

Recall that localization induces an injective map  $\text{Spec}(S^{-1}A) \rightarrow \text{Spec}^{-1}(A)$  with image  $\mathfrak{q}$  such that  $\mathfrak{q} \cap S = \emptyset$ . If  $S = A \setminus \mathfrak{p}$ , then this simplifies to  $\text{Spec}(A_{\mathfrak{p}}) = \mathfrak{q} \subseteq A$  such that  $\mathfrak{q} \subseteq \mathfrak{p}$ . In particular,  $A_{\mathfrak{p}}$  is a local ring with a unique maximal ideal.

**Proposition 2.10.** Let  $M$  be an  $A$ -module. Then, the following are equivalent:

- (i)  $M = 0$
- (ii) For all prime ideals  $\mathfrak{p}$ , we have  $M_{\mathfrak{p}} = 0$
- (iii) For all maximal ideals  $\mathfrak{m}$ , we have  $M_{\mathfrak{m}} = 0$



*Proof.* (i) implies (ii) implies (iii) is obvious.

To prove (iii) implies (i), suppose  $M \neq 0$ . Choose some non-zero  $x \in M$ . Then,  $\text{Ann}(x)$ , which denotes all  $a \in A$  such that  $ax = 0$  (recall that this is called the annihilator of  $x$ ), is a proper subset of  $A$ . Hence, there exists a maximal ideal  $\mathfrak{m}$  with  $\text{Ann}(x) \subseteq \mathfrak{m}$ .

Now, consider  $\frac{x}{1} \in M_{\mathfrak{m}}$ . If  $\frac{x}{1} = 0$  in  $M_{\mathfrak{m}}$ , then  $sx = 0$  for some  $s \in A \setminus \mathfrak{m}$ . However,  $(A \setminus \mathfrak{m}) \cap \text{Ann}(x) = \emptyset$ . Thus,  $\frac{x}{1} \neq 0$  in  $M_{\mathfrak{m}}$ , implying that  $M_{\mathfrak{m}} \neq 0$ .  $\square$

**Proposition 2.11.** Let  $f : M \rightarrow N$  be any a  $A$ -module homomorphism. Then, the following are equivalent:

- (i)  $f$  is injective
- (ii) For all prime ideals  $\mathfrak{p}$ ,  $f_{\mathfrak{p}} : M_{\mathfrak{p}} \rightarrow N_{\mathfrak{p}}$  is injective
- (iii) For all maximal ideals  $\mathfrak{m}$ ,  $f_{\mathfrak{m}} : M_{\mathfrak{m}} \rightarrow N_{\mathfrak{m}}$  is injective

### 2.3 Tensor Products

**Definition 2.8 (bilinear map).** Fix a ring  $A$ . Let  $M, N, P$  be  $A$ -modules. A map  $b : M \times N \rightarrow P$  is said to be bilinear if the following properties hold:

- (i)  $b(m + m', n) = b(m, n) + b(m', n)$  and  $b(m, n + n') = b(m, n) + b(m, n')$
- (ii)  $b(am, n) = b(m, an) = a \cdot b(m, n)$

We let

$\text{Bil}_A(M \times N, P)$  denote the set of bilinear maps  $b : M \times N \rightarrow P$  over  $A$ .

**Lemma 2.2.** There exists an  $A$ -module  $M \otimes_A N$  together with an  $A$ -bilinear map

$$b^{\text{univ}} : M \times N \rightarrow M \otimes_A N$$

such that the induced map

$$\text{Hom}_A(M \otimes_A N, P) \rightarrow \text{Bil}_A(M \times N, P) \quad \text{where} \quad f \rightarrow f \circ b^{\text{univ}}$$

is an isomorphism for all  $P$ .

*Proof.* We discuss the construction of  $M \otimes_A N$  as a module. Let  $F$  be the free  $A$ -module generated by all pairs  $(m, n)$ . Let  $R \subseteq F$  be the  $A$ -submodule generated by all elements of the following forms:

- (i)  $(m_1 + m_2, n) - (m_1, n) - (m_2, n)$
- (ii)  $(m, n_1 + n_2) - (m, n_1) - (m, n_2)$
- (iii)  $(am, n) - a(m, n)$
- (iv)  $(m, an) - a(m, n)$

Set  $M \otimes_A N = F/R$ . Consider the map

$$F \rightarrow M \otimes_A N \quad \text{where} \quad (m, n) \mapsto (m \otimes n).$$

By construction,  $M \otimes_A N$  is spanned by the elements  $m \otimes n$  and these satisfy the following relations:

- (i)  $(m_1 + m_2) \otimes n = m_1 \otimes n + m_2 \otimes n$


(ii)  $m \otimes (n_1 + n_2) = m \otimes n_1 + m \otimes n_2$

(iii)  $(am) \otimes n = a(m \otimes n) = m \otimes (an)$

Thus the map  $b^{\text{univ}} : M \times N \rightarrow M \otimes_A N$  sending  $(m, n) \mapsto m \otimes n$  is bilinear. It is also clear that if  $f : M \otimes_A N \rightarrow P$  is any  $A$ -module map, the induced map  $f \circ b^{\text{univ}} : M \times N \rightarrow P$  is  $A$ -bilinear. Conversely, suppose that we have a bilinear map  $\mathcal{B} : M \times N \rightarrow P$ . Define an  $A$ -module map  $\tilde{\mathcal{B}} : F \rightarrow P$  defined by

$$\sum a_i (m_i, n_i) \mapsto \sum a_i \mathcal{B}(m_i, n_i).$$

By the definition of bilinearity, we have  $R \subseteq \ker \tilde{\mathcal{B}}$ . Thus,  $\tilde{\mathcal{B}}$  factors as



for some unique  $A$ -module map  $\beta$ . Finally, it is clear that  $\beta \circ b^{\text{univ}} = \mathcal{B}$  by construction. □

**Proposition 2.12.** Here are some nice properties of the tensor product.

(i)  $M \otimes_A A = M$

(ii)  $M \otimes_A N \cong N \otimes_A M$

(iii)  $(M_1 \oplus M_2) \otimes_A N \cong (M_1 \otimes_A N) \oplus (M_2 \otimes_A N)$

(iv)  $(M \otimes_A N) \otimes_A K \cong M \otimes_A (N \otimes_A K)$

Here is a generalisation of (iv). Suppose we are given two rings  $A$  and  $B$ . let  $M$  and  $N$  be  $A$ - and  $B$ -modules respectively and  $P$  be an  $(A, B)$ -module. Then,

$$M \otimes_A P \text{ is a } B\text{-module} \quad \text{and} \quad P \otimes_B N \text{ is an } A\text{-module.}$$

Moreover,

$$(M \otimes_A P) \otimes_B N \cong M \otimes_A (P \otimes_B N).$$

Suppose  $M$  is an  $A$ -module and  $\varphi : A \rightarrow B$  is a ring map, i.e.  $B$  is an  $A$ -algebra. Then,  $M \otimes_A B$  is canonically a  $B$ -module, i.e.

$$b \left( \sum m_i \otimes b_i \right) = \sum m_i \otimes (bb_i).$$

This is compatible with the obvious  $A$ -module structure because

$$\varphi(a) \cdot \sum m_i \otimes b_i = \sum m_i \otimes \varphi(a) b_i = \sum am_i \otimes b_i = a \left( \sum m_i \otimes b_i \right).$$

**Proposition 2.13.** Fix  $S \subseteq A$  and let  $M$  be an  $A$ -module. Then, there exists a canonical isomorphism of  $S^{-1}A$  modules, i.e.

$$S^{-1}A \otimes_A M \cong S^{-1}M.$$

*Proof.* The map

$$S^{-1}A \times M \rightarrow S^{-1}M \quad \text{where} \quad \left( \frac{a}{s}, m \right) = \frac{am}{s}$$

is  $A$ -bilinear so it induces a unique  $A$ -module homomorphism as follows:

$$f : S^{-1}A \otimes_A M \rightarrow S^{-1}M \quad \text{where} \quad \sum \frac{a_i}{s_i} \otimes m_i \mapsto \sum \frac{a_i m_i}{s_i}$$

which is obviously surjective as

$$\frac{m}{s} = f \left( \frac{1}{s} \otimes m \right).$$

It suffices to prove that  $f$  is injective. Let

$$\sum \frac{a_i}{s_i} \otimes m_i \in S^{-1}A \otimes_A M \quad \text{be an arbitrary element.}$$

Set

$$s = \prod s_i \quad \text{and} \quad t_i = \prod_{j \neq i} s_j.$$

Then,

$$\sum \frac{a_i}{s_i} \otimes m_i = \sum \frac{a_i t_i}{s} \otimes m_i = \sum \frac{1}{s} \otimes a_i t_i m_i = \frac{1}{s} \otimes \sum a_i t_i m_i.$$

Thus, all elements of  $S^{-1}A \otimes_A M$  can be written in the form  $\frac{1}{s} \otimes m$  where  $m \in M$ . Thus,

$$f \left( \frac{1}{s} \otimes m \right) = \frac{m}{s} = 0 \quad \text{implies} \quad \text{there exists some } t \in S \text{ such that } tm = 0.$$

But then

$$\frac{1}{s} \otimes m = \frac{1}{ts} \otimes tm = \frac{1}{ts} \otimes 0 = 0.$$

To summarise,  $f \left( \frac{1}{s} \otimes m \right) = 0$  implies  $\frac{1}{s} \otimes m = 0$ , so  $f$  is injective. □

**Proposition 2.14 (tensor-hom adjunction).** Let  $M, N, P$  be  $A$ -modules. Then,

$$\text{Hom}_A(M \otimes_A N, P) \cong \text{Hom}_A(M, \text{Hom}_A(N, P)).$$

*Proof.* We have

$$\text{Hom}_A(M \otimes_A N, P) = \text{Bil}_A(M \times N, P).$$

This has the following canonical isomorphism:

$$\text{Bil}_A(M \times N, P) \cong \text{Hom}(M, \text{Hom}_A(N, P)) \quad \text{where} \quad b \mapsto (M \rightarrow \text{Hom}_A(N, P) \text{ where } m \mapsto b(m, \cdot))$$

To see this, observe that a bilinear map  $b : M \times N \rightarrow P$  naturally induces a map  $M \rightarrow \text{Hom}_A(N, P)$  by sending  $m \in M$  to the function  $N \rightarrow P$  defined by  $n \mapsto b(m, n)$ . Since  $b$  is bilinear, then the map  $M \rightarrow \text{Hom}_A(N, P)$  is  $A$ -linear. Conversely, given a module homomorphism  $f : M \rightarrow \text{Hom}_A(N, P)$ , define a bilinear map  $b : M \times N \rightarrow P$  by setting  $b(m, n) = f(m)(n)$ . This is a bilinear map since  $f$  is linear in  $m$  and  $f(m)$  is linear in  $n$ . □

Suppose  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  is exact and  $N$  is an  $A$ -module. Is  $0 \rightarrow M' \otimes_A N \rightarrow M \otimes_A N \rightarrow M'' \otimes_A N$  still an exact sequence? The answer is no, in general, but we have something less strict as follows.

**Proposition 2.15.** If

$$M' \rightarrow M \rightarrow M'' \rightarrow 0 \text{ is exact then } M' \otimes N \rightarrow M \otimes N \rightarrow M'' \otimes N \rightarrow 0 \text{ is exact.}$$

In other words, the functor  $- \otimes N$  is *right exact*.

To prove this, we need the following lemma:

**Lemma 2.3.** The following hold:

- (i) A sequence  $M' \rightarrow M \rightarrow M'' \rightarrow 0$  is exact if and only if for all  $A$ -modules  $P$ , the sequence  $0 \rightarrow \text{Hom}_A(M'', P) \rightarrow \text{Hom}_A(M, P) \rightarrow \text{Hom}_A(M', P)$  of abelian groups of homomorphisms is exact
- (ii) A sequence  $0 \rightarrow M' \rightarrow M \rightarrow M''$  is exact if and only if for all  $A$ -modules  $P$ , the sequence  $0 \rightarrow \text{Hom}_A(P, M') \rightarrow \text{Hom}_A(P, M) \rightarrow \text{Hom}_A(P, M'')$  is exact

We now prove Proposition 2.15.

*Proof.* We can argue now as follows: Suppose  $M' \rightarrow M \rightarrow M'' \rightarrow 0$  is exact. Since 0 is on the right, we use (i) of Lemma 2.3. We are going to use a *funny* choice of  $P$ , in particular, we see that the sequence

$$0 \rightarrow \text{Hom}_A(M'', \text{Hom}_A(N, P)) \rightarrow \text{Hom}_A(M, \text{Hom}_A(N, P)) \rightarrow \text{Hom}_A(M', \text{Hom}_A(N, P))$$

is exact by (i) of Lemma 2.3. By the tensor-hom adjunction (Proposition 2.14) we get

$$0 \rightarrow \text{Hom}_A(M'' \otimes_A N, P) \rightarrow \text{Hom}_A(M \otimes_A N, P) \rightarrow \text{Hom}_A(M' \otimes_A N, P).$$

Since this is exact for all  $P$ , using the reverse of the first part of Proposition 2.15, we get

$$0 \rightarrow M' \otimes_A N \rightarrow M \otimes_A N \rightarrow M'' \otimes_A N$$

which is exact. □

**Definition 2.9 (flat module).** An  $A$ -module  $N$  is flat if  $- \otimes_A N$  is exact. Equivalently, for every injective  $A$ -module map  $M' \rightarrow M$ , the induced map  $M' \otimes N \rightarrow M \otimes N$  is still injective.

**Example 2.17.** Let  $A$  be a commutative ring and  $S \subseteq A$  be a multiplicatively closed subset. Recall that the localization  $S^{-1}A$  consists of elements of the form  $\frac{a}{s}$ , where  $a \in A$  and  $s \in S$ . Here, the multiplication and addition operations are defined naturally.

$S^{-1}A$  is naturally an  $A$ -module via the action

$$a \cdot \frac{b}{s} = \frac{ab}{s} \quad \text{for all } a, b \in A, s \in S.$$

Next, for any  $A$ -module  $M$ , tensoring with  $S^{-1}A$  defines a functor

$$M \mapsto M \otimes_A S^{-1}A.$$

By the universal property of localization, we have

$$M \otimes_A S^{-1}A \cong S^{-1}M.$$

The map  $M \mapsto S^{-1}M$  is an exact functor, i.e. if we have an exact sequence of  $A$ -modules

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0,$$

then applying  $S^{-1}(-)$  yields an exact sequence of localized modules as follows:

$$0 \rightarrow S^{-1}M' \rightarrow S^{-1}M \rightarrow S^{-1}M'' \rightarrow 0$$

since localization commutes with taking kernels and cokernels. As  $- \otimes_A N$  is exact, then  $- \otimes_A S^{-1}A$  is also exact, making  $S^{-1}A$  a flat  $A$ -module.

**Example 2.18.** Let  $A$  be a ring. Suppose we are given ideals  $I, J \subseteq A$ . Then,

$$0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0 \quad \text{is exact.}$$

This is because  $I \rightarrow A$  is just the inclusion map, which is injective, and  $A \rightarrow A/I$  is the natural quotient map, which is surjective. Also, the kernel of the quotient map is  $I$ , so the sequence is exact.

We then tensor this sequence with  $A/J$ , meaning we apply  $- \otimes_A A/J$  to every term to obtain

$$I \otimes_A A/J \xrightarrow{\alpha} A/J \rightarrow A/I \otimes_A A/J \rightarrow 0 \quad \text{which is exact.}$$

To see why, for any element in  $I \otimes_A A/J$ , we have

$$\alpha \left( \sum i_j \otimes \bar{a}_j \right) = \sum \alpha (a_j i_j \otimes 1) = \sum \overline{a_j i_j}.$$

This means that elements in  $I \otimes_A A/J$  get mapped to their products modulo  $J$ .  $\text{im } \alpha$  consists of all elements in  $A/J$  that come from sums of products of elements from  $I$  and arbitrary elements of  $A$ . A crucial observation by the second isomorphism theorem yields

$$\text{im } \alpha = I / (I \cap J) \cong (I + J) / J.$$

Here,  $\text{coker } \alpha = A / (I + J)$ . Thus, we obtain the new exact sequence

$$I \otimes_A A/J \rightarrow A/J \rightarrow A / (I + J) \rightarrow 0.$$

Note that if the sequence is exact at  $I \otimes_A A/J$ , then  $\alpha$  is injective.

However, the sequence may not be flat! For example, let  $A = \mathbb{Z}$ ,  $I = (10)$  and  $J = (5)$ , so the sequence we tensor is

$$0 \rightarrow (10) \rightarrow \mathbb{Z} \rightarrow \mathbb{Z}/10\mathbb{Z} \rightarrow 0.$$

Tensoring with  $\mathbb{Z}/5\mathbb{Z}$ , we obtain

$$(10) \otimes_{\mathbb{Z}} \mathbb{Z}/5\mathbb{Z} \rightarrow \mathbb{Z}/5\mathbb{Z} \rightarrow \mathbb{Z}/5\mathbb{Z} \rightarrow 0.$$

Here,  $\text{im } \alpha$  consists of elements of the form  $10k \otimes \bar{1} \mapsto \overline{10k}$  in  $\mathbb{Z}/5\mathbb{Z}$ , which is in fact 0. Since every element in  $(10) \otimes_{\mathbb{Z}} \mathbb{Z}/5\mathbb{Z}$  is mapped to zero, then  $\text{im } \alpha = \{0\}$ , so  $\alpha$  is not injective, i.e. the sequence is not exact. This makes  $\mathbb{Z}/5\mathbb{Z}$  not a flat  $\mathbb{Z}$ -module.

Suppose we are given ring homomorphisms  $A \rightarrow B$  and  $A \rightarrow C$ . Then,  $B \otimes_A C$  is canonically equipped with a ring structure and satisfies a universal property. To see why, we shall explicitly construct the tensor product.

Consider the map


$$B \times C \times B \times C \rightarrow B \otimes_A C \quad \text{where} \quad (b, c, b', c') \mapsto bb' \otimes cc'.$$

This is a well-defined map of sets. Observe that this map is  $A$ -linear in each variable. We are going to spam some universal properties (of bilinear maps). We obtain the linear map  $(B \otimes_A C) \otimes_A (B \otimes_A C) \rightarrow B \otimes_A C$  such that the following diagram commutes:

$$\begin{array}{ccc} (B \otimes_A C) \otimes_A (B \otimes_A C) & \longrightarrow & B \otimes_A C \\ \uparrow & \nearrow \mu & \\ (B \otimes_A C) \times (B \otimes_A C) & & \end{array}$$

Composing with the universal map from the universal property of tensors, we obtain the multiplication map.

**Proposition 2.16 (universal property of pushout).** Given any commutative diagram of  $A$ -algebras



there exists a unique ring homomorphism  $h : B \otimes_A C \rightarrow D$  making the above diagram commute.

The idea in Proposition 2.16 is that given  $f$  and  $g$ , we can consider the map

$$B \times C \rightarrow D \quad \text{where} \quad (b, c) \mapsto f(b)g(c)$$

This is well-defined and  $A$ -bilinear (since we are working with  $A$ -algebras), thus the universal property for bilinear maps yields the desired map  $B \otimes_A C \rightarrow D$ .

**Example 2.19.** Fix a ring  $A$  and consider ring homomorphisms  $A \rightarrow B$  and  $A \rightarrow A[X]$ . Note that  $B \otimes_A A[X] = B[X]$ . Also, the following diagram commutes:

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ A[X] & \longrightarrow & B \otimes_A A[X] = B[X] \end{array}$$

To deduce this without excessive calculations, we can simply write

$$A[X] = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} A$$

and use the fact that tensor products distribute across direct sums. Alternatively, we can use the universal property from before.

**Example 2.20.** Say we are interested in  $\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C}$ . Recall from MA3201 that  $\mathbb{C} = \mathbb{R}[X]/(X^2 + 1)$ . Then using a previous result that  $(A/J) \otimes_A B = B/JB$  (where  $JB$  is the ideal in  $B$  generated by the image of  $J$ ), we obtain

$$\begin{aligned} \mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} &= \mathbb{R}[X]/(X^2 + 1) \otimes_{\mathbb{R}} \mathbb{C} \\ &= (\mathbb{R}[X] \otimes_{\mathbb{R}} \mathbb{C}) / (X^2 + 1) \\ &= \mathbb{C}[X] / (X^2 + 1) \\ &= \mathbb{C}[X] / ((X + i)(X - i)) \\ &= \mathbb{C} \times \mathbb{C} \end{aligned}$$

We can define an explicit isomorphism

$$\mathbb{C} \otimes_{\mathbb{R}} \mathbb{C} \cong \mathbb{C} \times \mathbb{C} \quad \text{where} \quad a \otimes b \mapsto (\bar{a}b, ab).$$

In fact, the same proof will work to prove that for any finite Galois extension  $E/F$ , we have  $E \otimes_F E \cong E^{[E:F]}$ .

# Chapter 3

## Some Classes of Rings

### 3.1 Noetherian Rings

**Definition 3.1 (Noetherian ring).** Given a ring  $A$ , the following are equivalent:

- (i) Every ascending chain of ideals  $I_1 \subseteq I_2 \subseteq \dots$  stabilizes, i.e.  $I_n = I_{n+1}$  for sufficiently large  $n \in \mathbb{N}$
  - (ii) Every ideal  $I \subseteq A$  is finitely generated as an  $A$ -module.
  - (iii) For any  $N \subseteq M$  of  $A$ -modules,  $M$  is finitely generated implies  $N$  is finitely generated
- If a ring satisfies any (thus all) of these conditions, we say that the ring is Noetherian.

We can also see Definition 3.1 as a proposition. We now provide a proof for it.

*Proof.* We first prove (i) implies (ii) by contraposition. Assume that  $I \subseteq A$  is an ideal which is not finitely generated as an  $A$ -module. Then, we can pick generators  $I = (a_1, a_2, \dots)$  such that  $(a_1) \subsetneq (a_1, a_2) \subsetneq \dots$ . This gives a chain  $I_1 \subsetneq I_2 \subsetneq I_3 \subsetneq \dots$  that does not stabilize. Therefore (i) does not hold. As such, (i) implies (ii).

We then prove (ii) implies (iii) by performing induction on the minimum number of generators of  $M$ . If  $M$  is generated by one element, then  $M$  is isomorphic to a quotient  $A/I$  of  $A$  for some ideal  $I$ . Then submodules of  $M$  are identified with ideals  $J$  of  $A$  that contain  $I$ . Since all ideals of  $A$  are finitely generated by assumption,  $J$  is finitely generated, whence  $J/I \cong N$  is finitely generated.

For the induction hypothesis, suppose we are given an  $A$ -module  $M$  generated by  $x_1, \dots, x_n$ . Let  $M'$  be a submodule generated by  $x_1$ , let  $M'' = M/M'$ . This yields a short exact sequence as follows:

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

where since  $M'$  is generated by  $x_1$ ,  $M''$  must be generated by (the image in  $M''$  of)  $x_2, \dots, x_n$ . Now we pick any submodule  $N \subset M$ . Then, we obtain the following diagram:

$$\begin{array}{ccccccccc}
 0 & \longrightarrow & M' & \longrightarrow & M & \longrightarrow & M'' & \longrightarrow & 0 \\
 & & \uparrow & & \uparrow & & \uparrow & & \\
 0 & \longrightarrow & N \cap M' & \longrightarrow & N & \longrightarrow & N/(N \cap M') & \longrightarrow & 0 \\
 & & & & & & \parallel & & \\
 & & & & & & (N + M')/M' & & 
 \end{array}$$

Then,  $N \cap M'$  is finitely generated by the base case (the case of modules generated by one element) and  $N/(N \cap M')$  is finitely generated by the induction hypothesis. Therefore,  $N$  is also finitely generated by Example 2.7<sup>†</sup>.

Lastly, we prove (iii) implies (i). Say any submodule of a finitely generated  $A$ -module is finitely generated,

<sup>†</sup>In Example 2.7, recall we proved that if the outer two modules are finitely generated, then the middle module is also finitely generated.



then given any ascending chain of ideals  $I_1 \subseteq I_2 \subseteq \dots$ , the union

$$\bigcup_{n \in \mathbb{N}} I_n \subseteq A \text{ is finitely generated as an } A\text{-module.}$$

Thus,

$$\bigcup_{n \in \mathbb{N}} I_n = (x_1, \dots, x_j) \text{ for some } x_1, \dots, x_j \in I_N,$$

where  $N \in \mathbb{N}$  is sufficiently large. It follows that  $I_n = I_{n+1}$  for all  $n \geq N$ . □

**Example 3.1.**  $\mathbb{Z}$  is a Noetherian ring.

**Example 3.2.** Any field is a Noetherian ring.

**Example 3.3.** If  $A$  is a Noetherian ring and  $I \subseteq A$  is an ideal, then  $A/I$  is Noetherian.

**Example 3.4.** If  $A$  is Noetherian and  $S \subseteq A$  is multiplicatively closed, then  $S^{-1}A$  is Noetherian. To see why, any ideal  $I \subseteq S^{-1}A$  is extended from some  $I' \subseteq A$ . If  $x_1, \dots, x_n$  generate  $I'$ , then  $\frac{x_1}{1}, \dots, \frac{x_n}{1}$  generate  $I' (S^{-1}A) = I$ .


**Theorem 3.1 (Hilbert basis theorem).** If  $A$  is a Noetherian ring, then  $A[X]$  is also Noetherian.

*Proof.* See my MA4273 notes. □

**Corollary 3.1.** If  $A$  is Noetherian, then  $A[X_1, \dots, X_n]$  is also Noetherian.

**Definition 3.2 (finitely generated and finitely presented algebras).** Fix a ring  $A$ . Let  $B$  be an  $A$ -algebra. Then,  $B$  is *finitely generated* (or of *finite type*) as an  $A$ -algebra if and only if there exist ring homomorphisms such that the following diagram commutes, i.e. there exists

an isomorphism  $B \cong A[X_1, \dots, X_n]/I$  for some ideal  $I$  of  $A[X_1, \dots, X_n]$ .



Moreover, if there exists such a diagram where  $I$  is finitely generated as a  $A[X_1, \dots, X_n]$ -module, then we say that  $B$  is a *finitely presented*  $A$ -algebra.

**Proposition 3.1.** If  $A$  is Noetherian, then any finitely generated  $A$ -algebra  $B$  is Noetherian and finitely presented.

### 3.2

## A Preview of Dimension Theory

**Definition 3.3 (height of prime ideal).** Given a prime ideal of a ring  $\mathfrak{p} \subseteq A$ , define

$$\text{ht}(\mathfrak{p}) = \sup \{n \mid \text{there exists a chain } \mathfrak{p}_0 \subseteq \mathfrak{p}_1 \subseteq \dots \subseteq \mathfrak{p}_n = \mathfrak{p}\}$$

to be the height of  $\mathfrak{p}$ .

**Definition 3.4 (Krull dimension).** Given a prime ideal of a ring  $\mathfrak{p} \subseteq A$ , define

$$\dim(A) = \sup_{\mathfrak{p}} \text{ht}(\mathfrak{p}) = \sup \{n \mid \text{there exists a chain } \mathfrak{p}_0 \subsetneq \mathfrak{p}_1 \subsetneq \dots \subsetneq \mathfrak{p}_n\}.$$

**Example 3.5 (Krull dimension of a field).** For any field  $k$ , we have  $\dim(k) = 0$ . This is easy to see — the only prime ideals of  $k$  are itself and the zero ideal  $(0)$ . However,  $k$  is not a proper ideal of itself. As the longest possible chain of prime ideals is  $(0)$ , and there are no prime ideals strictly contained in  $(0)$  and no proper prime ideals strictly containing  $(0)$ , the longest chain of prime ideals has length 0.

**Example 3.6.**  $\dim(\mathbb{Z}) = 1$  and  $\text{ht}((p)) = 1$  for all primes  $p$ .

**Example 3.7.** For any field  $k$ ,  $\dim(k[X_1, \dots, X_n]) \geq n$  because  $0 \subseteq (X_1) \subseteq \dots \subseteq (X_1, \dots, X_n)$ .

**Theorem 3.2.** If  $A$  is Noetherian, then  $\dim(A[X]) = \dim(A) + 1$ .

Even if  $A$  is a Noetherian ring, its Krull dimension can still be infinite. An example of a Noetherian ring with infinite Krull dimension was constructed by Nagata in the 1950s. For the interested reader, please refer to Exercise 9.6 of ‘Commutative Algebra’ by D. Eisenbud.

**Proposition 3.2.**  $\text{ht}(\mathfrak{p}) = \dim(A_{\mathfrak{p}})$

**Theorem 3.3.**  $\dim(A_{\mathfrak{p}})$  is finite for any Noetherian ring.

**Remark 3.1.**  $\text{ht}(\mathfrak{p}) = \dim(A_{\mathfrak{p}})$ .

**Theorem 3.4.**  $\dim(A_{\mathfrak{p}})$  is finite for any Noetherian ring.

Here is a fun fact: suppose that  $A$  is not Noetherian but  $\dim A$  is finite. Then  $\dim(A) + 1 \leq \dim(A[X]) \leq 2\dim(A)$ , and there exist examples exhibiting every possibility in this range.

### 3.3

#### Integral Dependence and Integral Rings

**Definition 3.5.** Suppose  $A \subseteq B$ . An element  $x \in B$  is said to be *integral* over  $A$  if  $x$  satisfies some monic polynomial with coefficients in  $A$ , i.e. if

$$x^n + a_{n-1}x^{n-1} + \dots + a_1x + a_0 = 0 \quad \text{for some } a_0, \dots, a_{n-1} \in A.$$

**Example 3.8.** If  $A = \mathbb{Z}$  and  $B = \mathbb{Q}$ , then the set of elements in  $\mathbb{Q}$  integral over  $\mathbb{Z}$  is just  $\mathbb{Z}$ .

**Definition 3.6.** A module  $M$  is *faithful* if and only if  $\text{Ann}(M) = 0$ , i.e.  $R \rightarrow \text{Hom}_R(M, M)$  is injective.

**Example 3.9.**  $\mathbb{Z}$  is a faithful  $\mathbb{Z}$ -module. To see why, note that

$$\text{Ann}(\mathbb{Z}) = \{n \in \mathbb{Z} : n\mathbb{Z} = 0\},$$

which forces  $n = 0$ , so  $\text{Ann}(\mathbb{Z}) = 0$ .

Note that  $A[x]$  is not a polynomial ring, but rather the submodule of  $B$  generated by  $A$ -multiples of powers of  $x$ . Thus  $A[x]$  consists of elements of  $B$  of the form

$$\sum_i a_i x^i.$$

**Proposition 3.3.** Fix  $A \subseteq B$ . Then, fix  $x \in B$ . The following are equivalent:

- (i)  $x$  is integral over  $A$
- (ii)  $A[x]$  is a finitely generated  $A$ -module
- (iii) There exists a finitely generated  $B$ -module  $C$  such that  $A[x] \subseteq C \subseteq B$
- (iv) There exists a faithful  $A[x]$ -module  $M$  which is finitely generated as an  $A$ -module

*Proof.* (i) implies (ii) is trivial. To see why, if  $x$  satisfies  $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$ , then  $A[x]$  is generated by  $1, x, x^2, \dots, x^{n-1}$  since all combinations of powers of  $x$  greater than  $n - 1$  can be reduced to lower terms by the relation.

(ii) implies (iii) by taking  $C = A[x]$ ; (iii) implies (iv) by taking  $M = C$ .

Lastly, to prove (iv) implies (i), suppose we are given a finitely generated  $A$ -module  $M$  equipped with a faithful action by the elements of  $A[x]$ . Consider the  $A$ -module endomorphism  $\phi : M \rightarrow M$  where  $m \mapsto xm$ . By applying the useless version of Nakayama's lemma (Proposition 2.2) to the whole ring  $R = J = A$  and  $M = N$ , and  $\phi$  as defined previously, we then get, as an  $A$ -module endomorphism on  $M$ ,

$$x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$$

Since  $M$  is a faithful  $A[x]$ -module, then  $A[x] \hookrightarrow \text{Hom}_A(M, M)$  is injective, so that  $x^n + a_{n-1}x^{n-1} + \dots + a_0 = 0$  in  $A[x]$ . The result follows.  $\square$

**Corollary 3.2.** Given  $A \subseteq B$  and  $x_1, \dots, x_n \in B$  each integral over  $A$ , then  $A[x_1, \dots, x_n] \subseteq B$  is a finitely generated  $A$ -module.

**Definition 3.7.** Let  $f : A \rightarrow B$  be a ring homomorphism. We say that it is *integral* if and only if  $f(A) \subseteq B$  is an integral ring extension, i.e.  $B$  is an integral ring extension of  $f(A)$ .

**Proposition 3.4 (transitivity of integral ring extension).** Given  $A \subseteq B \subseteq C$  are ring extensions, and  $A \subseteq B$  and  $B \subseteq C$  are integral ring extensions, then  $A \subseteq C$  is an integral ring extension.

*Proof.* Pick  $x \in C$ . Then, there exists an equation

$$x^n + b_1x^{n-1} + \dots + b_n = 0 \quad \text{where } b_i \in B.$$

Since  $B$  is integral over  $A$ , then  $B' = A[b_1, \dots, b_n] \subseteq C$  is a finitely generated  $A$ -module. So,  $B'[x]$  is also a finitely generated  $B'$ -module, so it is finitely generated as an  $A$ -module. Thus  $A[x] \subseteq B'[x] \subseteq C$  with  $B'[x]$  finitely generated as an  $A$ -module. To complete the proof, recall that (iii) implies (i) in Proposition 3.3.  $\square$

**Corollary 3.3.** Let  $A \subseteq B$  be a ring extension and  $C$  be the integral closure of  $A$  in  $B$ . Then,  $C$  is also integrally closed in  $B$ .

*Proof.* If  $x \in B$  is integral over  $C$ , then  $A \subseteq C \subseteq C[x]$  are integral ring extensions. So,  $A \subseteq C[x]$  is an integral ring extension. So,  $x$  is integral over  $A$ . We conclude that  $x \in C$ .  $\square$

**Proposition 3.5.** Let  $A \subseteq B$  be an integral ring extension. Then, the following hold:

(i) Let  $J \subseteq B$  be an ideal. Then,

$$A/(J \cap A) = B/J \quad \text{is an integral ring extension.}$$

(ii) Let  $S \subseteq A$  be a multiplicatively closed subset. Then,

$$S^{-1}A \subseteq S^{-1}B \quad \text{is an integral ring extension.}$$

*Proof.* For (i), suppose we are given  $b \in B/J$  and a lift  $\tilde{b} \in B$ . Note that  $\tilde{b}$  satisfies

$$\tilde{b}^n + a_1 \tilde{b}^{n-1} + \dots + a_n = 0 \quad \text{where } a_i \in A.$$

Then, modulo  $J \cap A$ , we obtain

$$b^n + \bar{a}_1 b^{n-1} + \dots + \bar{a}_n = 0 \quad \text{where } \bar{a}_i \in A/(J \cap A).$$

This proves (i). As for (ii), pick any  $\frac{x}{s} \in S^{-1}B$ . Then, we can pick

$$x^n + a_1 x^{n-1} + \dots + a_n = 0 \quad \text{in } B.$$

As such, we obtain

$$\frac{x^n}{s^n} + \frac{a_1}{s} \cdot \frac{x^{n-1}}{s^{n-1}} + \dots + \frac{a_n}{s^n} = 0 \quad \text{in } S^{-1}B.$$

Thus,  $t = \frac{x}{s}$  is a root of the polynomial equation

$$t^n + \frac{a_1}{s} t^{n-1} + \dots + \frac{a_n}{s^n} = 0,$$

which proves (ii).  $\square$

**Proposition 3.6.** Let  $A \subseteq B$  be an integral ring extension with both rings being integral domains. Then,

$$A \text{ is a field} \quad \text{if and only if} \quad B \text{ is a field.}$$

*Proof.* We first prove the forward direction. Suppose  $A$  is a field. Let  $0 \neq x \in B$  be an element. Then, there exists an equation

$$x^n + a_1 x^{n-1} + \dots + a_{n-1} x + a_n = 0 \quad \text{where } a_i \in A.$$

Note that  $a_n \neq 0$  since  $A$  is an integral domain. Then,

$$y = -a_n^{-1} (x^{n-1} + a_1 x^{n-2} + \dots + a_{n-1}) \text{ in } B \quad \text{satisfies the equation } xy = 1.$$

As such,  $B$  is a field.

For the reverse direction, suppose  $B$  is a field. Choose  $0 \neq x \in A$ . Then,  $x^{-1} \in B$  is integral over  $A$  so we obtain the equation

$$x^{-n} + a_1 x^{-n+1} + \dots + a_n = 0 \quad \text{where } a_i \in A.$$

Then,

$$x^{-1} = - (a_1 + a_2 x + a_3 x^2 + \dots + a_n x^{n-1}) \in A.$$

This implies that  $x^{-1} \in A$ , so  $A$  is a field.  $\square$

**Corollary 3.4.** Let  $A \subseteq B$  be an integral ring extension and  $\mathfrak{q} \subseteq B$  be a prime ideal. Let  $\mathfrak{p} = A \cap \mathfrak{q}$  be its contraction. Then,

$$\mathfrak{q} \text{ is maximal if and only if } A/\mathfrak{p} \text{ is a field.}$$

*Proof.* Note that  $A/\mathfrak{p} \subseteq B/\mathfrak{q}$  is an integral ring extension, where  $A$  and  $B$  are integral domains. Recall that

$$\mathfrak{p} \text{ is a maximal ideal if and only if } A/\mathfrak{p} \text{ is a field.}$$

Hence, this is equivalent to saying that  $B/\mathfrak{q}$  is a field and  $\mathfrak{q}$  is a maximal ideal. □

**Proposition 3.7.** Let  $A \subseteq B$  be an integral ring extension. Suppose we are given

$$\mathfrak{q} \subseteq \mathfrak{q}' \subseteq B \text{ as prime ideals such that } \mathfrak{q} \cap A = \mathfrak{q}' \cap A = \mathfrak{p}.$$

Then,  $\mathfrak{q} = \mathfrak{q}'$ .

*Proof.* It is clear that  $\mathfrak{q}$  and  $\mathfrak{q}'$  both contain the extension of  $\mathfrak{p}$ . Set  $A_{\mathfrak{p}}$  as usual and  $B_{\mathfrak{p}} = (A \setminus \mathfrak{p})^{-1}B$ . So,  $A_{\mathfrak{p}} \subseteq B_{\mathfrak{p}}$  is an integral ring extension. Let  $\mathfrak{m} = \mathfrak{p}A_{\mathfrak{p}}$ , and  $\mathfrak{n} = \mathfrak{q}B_{\mathfrak{p}}$ ,  $\mathfrak{n}' = \mathfrak{q}'B_{\mathfrak{p}}$ .

By construction,  $\mathfrak{n}$  and  $\mathfrak{n}'$  contract to  $\mathfrak{q}$  and  $\mathfrak{q}'$  respectively in  $B$ . But contracting  $\mathfrak{n}$  and  $\mathfrak{n}'$  in  $A_{\mathfrak{p}}$  yields  $\mathfrak{m}$ , and  $A_{\mathfrak{p}} \subseteq B_{\mathfrak{p}}$  is integral, so  $\mathfrak{n} \subseteq \mathfrak{n}'$  are both maximal. □

**Theorem 3.5.** Let  $A \subseteq B$  be an integral ring extension and let  $\mathfrak{p} \subseteq A$  be a prime ideal. Then there exists a prime ideal  $\mathfrak{q} \subseteq B$  with  $A \cap \mathfrak{q} = \mathfrak{p}$ . In other words, there exists a surjective map  $\text{Spec}(B) \rightarrow \text{Spec}(A)$  (given by contraction).

*Proof.* Consider the following diagram of ring homomorphisms:

$$\begin{array}{ccc} A & \hookrightarrow & B \\ \downarrow & & \downarrow \\ A_{\mathfrak{p}} & \hookrightarrow & B_{\mathfrak{p}} \end{array}$$

Here, the horizontal arrows are integral extensions. Pick any maximal ideal  $\mathfrak{n}$  in  $B_{\mathfrak{p}}$ . Then  $\mathfrak{n} \cap A_{\mathfrak{p}}$  is maximal in the local ring  $A_{\mathfrak{p}}$ , so  $\mathfrak{n} \cap A_{\mathfrak{p}} = \mathfrak{p}A_{\mathfrak{p}}$ . Thus,  $\mathfrak{n} \cap A = \mathfrak{p}$  (this is contracted into  $A$ ). Then  $\mathfrak{q} := \mathfrak{n} \cap B$  is prime and  $\mathfrak{q} \cap A = \mathfrak{p}$ , as desired. □

We then discuss a *funny theorem* known as the *going up theorem*. This is also known as the Cohen–Seidenberg theorem.

**Theorem 3.6 (Cohen–Seidenberg theorem, going up).** Let  $A \subseteq B$  be an integral ring extension and

$$\begin{aligned} \mathfrak{p}_1 \subseteq \mathfrak{p}_2 \subseteq \dots \subseteq \mathfrak{p}_n & \text{ be a chain of prime ideals in } A \\ \mathfrak{q}_1 \subseteq \mathfrak{q}_2 \subseteq \dots \subseteq \mathfrak{q}_m & \text{ be a chain of prime ideals in } B \end{aligned}$$

where  $m \leq n$  such that  $\mathfrak{q}_i \cap A = \mathfrak{p}_i$  for all  $1 \leq i \leq m$ . Then, we can extend the chain to  $\mathfrak{q}_1 \subseteq \dots \subseteq \mathfrak{q}_n$  such that  $\mathfrak{q}_i \cap A = \mathfrak{p}_i$  for all  $1 \leq i \leq n$ .

*Proof.* It suffices to consider where we can lengthen the chain of  $\mathfrak{q}$ 's one step at a time. This reduces to the case  $n = 2$  and  $m = 1$ .

Set  $\bar{A} = A/\mathfrak{p}_1 \subseteq \bar{B} = B/\mathfrak{q}$ . This is an integral ring extension. Since  $\mathfrak{p}_2$  contains  $\mathfrak{p}_1$ ,  $\bar{\mathfrak{p}}_2 = \mathfrak{p}_2/\mathfrak{p}_1 \subseteq \bar{A}$  is a prime ideal, and thus there exists  $\mathfrak{q}' \subseteq \bar{B}$  with  $\mathfrak{p}' \cap \bar{A} = \bar{\mathfrak{p}}_2$ . Then,  $\mathfrak{q}_2$ , defined as the preimage of  $\mathfrak{q}'$  along  $B \rightarrow \bar{B}$ , has the desired property.  $\square$

**Corollary 3.5.** If  $A \subseteq B$  is an integral ring extension, then  $\dim A \leq \dim B$ .

*Proof.* This is immediate from the going up theorem (Theorem 3.6) since given any chain of prime ideals in  $A$ , we can find a corresponding chain of prime ideals in  $B$  that is at least as long.  $\square$

**Theorem 3.7 (Cohen–Seidenberg theorem, going down).** Let  $A \subseteq B$  be an integral ring extension and

$$\begin{aligned} \mathfrak{p}_1 \supseteq \mathfrak{p}_2 \supseteq \dots \supseteq \mathfrak{p}_n & \text{ be a chain of prime ideals in } A \\ \mathfrak{q}_1 \supseteq \mathfrak{q}_2 \supseteq \dots \supseteq \mathfrak{q}_m & \text{ be a chain of prime ideals in } B \end{aligned}$$

where  $m \leq n$  such that  $\mathfrak{q}_i \cap A = \mathfrak{p}_i$  for all  $1 \leq i \leq m$ . Then, we can extend the chain to  $\mathfrak{q}_1 \supseteq \dots \supseteq \mathfrak{q}_n$  such that  $\mathfrak{q}_i \cap A = \mathfrak{p}_i$  for all  $1 \leq i \leq n$ .

**Proposition 3.8.** Suppose we are given a ring extension  $A \subseteq B$  (not necessarily integral), and  $C$  is the integral closure of  $A$  in  $B$ . Let  $S \subseteq A$  be any multiplicatively closed subset. Then,

$$S^{-1}C \text{ is the integral closure of } S^{-1}A \text{ in } S^{-1}B.$$

*Proof.* We already know that  $S^{-1}C$  is integral over  $S^{-1}A$ . Now, suppose  $\frac{b}{s} \in S^{-1}B$  is integral over  $S^{-1}A$ , i.e. there exists an equation:

$$\left(\frac{b}{s}\right)^n + \frac{a_1}{s_1} \left(\frac{b}{s}\right)^{n-1} + \dots + \frac{a_n}{s_n} = 0$$

with  $\frac{a_i}{s_i} \in S^{-1}A$ . Set  $t = s_1 \dots s_n$  and multiply the previous equation by  $(st)^n$ . After some rearrangement, we see that  $bt \in B$  is integral over  $A$ , so  $\frac{b}{s} = \frac{bt}{st} \in S^{-1}C$ .  $\square$

**Definition 3.8 (field of fractions).** Let  $A$  be an integral domain. Then,

$$\text{Frac}(A) = (A \setminus \{0\})^{-1}A \text{ is a field.}$$

We call this the field of fractions of  $A$ .

**Definition 3.9 (integrally closed domain).** A domain  $A$  is said to be an integrally closed domain if  $A$  is integrally closed in  $\text{Frac}(A)$ .

**Example 3.10.**  $\mathbb{Z}$  is an integrally closed domain.

**Example 3.11.** However,  $\mathbb{Z}[\sqrt{5}]$  is not an integrally closed domain because its integral closure in  $\mathbb{Q}(\sqrt{5})$  is

$$\mathbb{Z}\left[\frac{1+\sqrt{5}}{2}\right].$$

**Theorem 3.8.** Let  $A$  be an integral domain. Then, the following are equivalent:

- (i)  $A$  is integrally closed

- (ii)  $A_p$  is integrally closed for all prime ideals  $p$
- (iii)  $A_m$  is integrally closed for all maximal ideals  $m$

**Definition 3.10.** A ring  $A$  is normal if  $A_p$  is an integrally closed domain for all prime ideals  $p$ .

### 3.4 Completions

**Definition 3.11 (directed set).** A directed set is a set  $\Lambda$  with a relation  $R$  which is reflexive and transitive and such that

$$\text{for all } a, b \in \Lambda \text{ we have } a \leq c \text{ and } b \leq c.$$

We have the following chain of inclusions:

$$\text{totally ordered sets} \subseteq \text{directed sets} \subseteq \text{partially ordered sets}$$

Now, fix a ring  $A$  and let  $M$  be an  $A$ -module. Suppose we are given a collection  $\{M_\lambda\}_{\lambda \in \Lambda}$  of submodules  $M_\lambda \subseteq M$ , where  $\Lambda$  is a directed set with  $M_\lambda$ 's ordered by inclusion. Given this data, we can put a topology on  $M$ .

We declare that the open sets of  $M$  are arbitrary unions of the cosets of the  $M_\lambda$ 's. In other words,

$$\{x + M_\lambda : x \in M, \lambda \in \Lambda\} \text{ forms a base for this topology.}$$

Note that this makes  $M$  into a topological Abelian group under addition because  $M \times M \xrightarrow{+} M$  has the property that the preimage of any  $x + M_\lambda \supseteq (x + M_\lambda) \times M_\lambda$ . Moreover,  $A$ -multiplication is continuous because  $a : M \times M$  has the property that the pre-image of  $ax + M_\lambda \supseteq x + M_\lambda$ .  $M$  is what we call a linearly topologized  $A$ -module.

**Example 3.12.** If  $M = A$ , then all  $M_\lambda$ 's are ideals, and one can check that the multiplication map  $\cdot$  is continuous because

$$(x + M_\lambda)(y + M_\lambda) = xy + M_\lambda.$$

In particular,  $A$  is a topological ring.

**Example 3.13 ( $I$ -adic topology on  $M$ ).** Let our directed set  $\Lambda = \mathbb{N}$ , and let  $I \subset A$  be some fixed ideal. For any  $A$ -module  $M$ , define  $M_n = I^n M$ , and let  $M_n = I^n M$  for any  $A$ -module  $M$ .

The  $I$ -adic topology need not be Hausdorff — in fact, it can be quite nasty in general. For example, in Example 3.13, let  $I = (0)$ , then our topology becomes the one-point set topology. One can check that the topology induced on  $M$  induced by  $\{M_\lambda\}_{\lambda \in \Lambda}$  is Hausdorff if and only if

$$\bigcap_{\lambda \in \Lambda} M_\lambda = \{0\}.$$

Consider  $M/M_\lambda$  with its quotient topology. As  $M_\lambda$  is open and closed (since its complement is just a union of cosets of it), it follows that the preimage of any  $S \subset M/M_\lambda$  is

$$\bigcup_{s \in \tilde{S}} s + M_\lambda \text{ where } s + M_\lambda \in \tilde{S} \text{ refers to the lift of } s \in S$$

is open and closed. In other words, the inherited quotient topology of  $M/M_\lambda$  is discrete.

**Definition 3.12** (separation quotient). Define

$$M^{\text{sep}} = M / \bigcap_{\lambda \in \Lambda} M_\lambda.$$

This set inherits an inherent linear topology coming from

$$\left\{ \left( M_\lambda + \bigcap_{\mu \in \Lambda} M_\mu \right) / \bigcap_{\mu \in \Lambda} M_\mu \right\}_{\lambda \in \Lambda} \text{ is Hausdorff.}$$

**Definition 3.13** (completion of module). We define

$$\widehat{M} = \varprojlim M/M_\lambda \text{ to be the completion of } M.$$

Observe that for all  $\lambda \leq \mu$ , we have canonical maps  $\varphi_{\lambda\mu} : M/M_\mu \rightarrow M/M_\lambda$  (since we can quotient the larger submodule by the smaller submodule) such that  $\varphi_{\lambda\mu} \circ \varphi_{\mu\gamma} = \varphi_{\lambda\gamma}$  for all  $\lambda \leq \mu \leq \gamma$ . By definition,

$$\widehat{M} = \left\{ \{x_\lambda\}_{\lambda \in \Lambda} \in \prod_{\lambda} M/M_\lambda \mid \varphi_{\lambda\mu}(x_\mu) = x_\lambda \text{ for all } \lambda \leq \mu \right\} \subseteq \prod_{\lambda \in \Lambda} M/M_\lambda.$$

Equip  $\widehat{M}$  the subspace topology for

$$\subseteq \prod_{\lambda \in \Lambda} M/M_\lambda,$$

where  $M/M_\lambda$  is in the discrete topology and the product of  $M/M_\lambda$  over all  $\lambda \in \Lambda$  is in the product topology. A crucial observation here is that the following diagram commutes:

$$\begin{array}{ccc} M & \xrightarrow{\psi} & \widehat{M} \\ & \searrow & \downarrow p_\lambda \\ & & M/M_\lambda \end{array}$$

Here,  $\psi(x) = \{x\}_{\lambda \in \Lambda}$  and  $p_\lambda$  is the obvious projection of  $\widehat{M}$  to the corresponding coordinate  $\lambda \in \Lambda$ . By the commutativity of the diagram and the obvious surjectivity of taking quotients (since  $p_\lambda$  is a projection map), we deduce that  $p_\lambda$  is also surjective.

We also note that

$$\ker \psi = \bigcap_{\lambda \in \Lambda} M_\lambda \quad \text{and} \quad \text{im } \psi = \psi(M) \subseteq \widehat{M} \text{ is dense in } M.$$

This is seen from the following proposition (Proposition 3.9).

**Proposition 3.9.**  $\widehat{M}$  is linearly topologized by the submodules  $M_\lambda^* = \ker p_\lambda$ , and  $\ker p_\lambda$  is equal to the closure of  $\psi(M_\lambda)$  in  $\widehat{M}$ . Moreover,  $\widehat{M}$  is complete, in the sense that  $\widehat{\widehat{M}} \cong \widehat{M}$  as topological  $A$ -modules.

**Example 3.14.** Let  $A$  be a ring  $x_1, \dots, x_n \in A$ ,  $I = (x_1, \dots, x_n)$ , then  $\{I^j M\}_{j \in \mathbb{N}}$  gives the  $I$ -adic topology. We could also consider the directed system of ideals  $J_n = (x_1^n, \dots, x_n^n)$ . We claim that the two directed sets of ideals induce the same topology. Indeed, it is clear that  $J_m \subseteq I^m$ . Also  $I^{mn} \subseteq J_m$ . Therefore these two systems define the same topology on any  $A$ -module.



Now suppose that  $M$  is topologized by  $\{M_\lambda\}_{\lambda \in \Lambda}$  and let  $N \subseteq M$  be a submodule. Observe that the closure of  $N$  in  $M$  is  $\bar{N} = \bigcap_{\lambda \in \Lambda} (N + M_\lambda)$ . Indeed, we have the following chain of equivalent statements:

$$\begin{aligned} x \in \bar{N} & \text{ if and only if } (x + M_\lambda) \cap N \neq \emptyset \text{ for all } \lambda \in \Lambda \\ & \text{ if and only if } x \in \bigcap_{\lambda \in \Lambda} (N - M_\lambda) = \bigcap_{\lambda \in \Lambda} (N + M_\lambda) \end{aligned}$$

**Proposition 3.10.** Let  $N \subseteq M$  be a submodule, and let  $M'_\lambda$  be the image of  $M_\lambda \rightarrow M \rightarrow M/N$ . Then the quotient topology on  $M/N$  is the same as the linear topology induced by  $\{M'_\lambda\}_{\lambda \in \Lambda}$ .

*Proof.* Note that  $\bar{U} \subseteq M/N$  is open in the quotient topology if and only if its preimage  $U$  is open on  $M$ , i.e. for all  $x \in U$ , there exists  $\lambda \in \Lambda$  satisfying the following chain of equivalent statements:

$$\begin{aligned} \bar{U} \subseteq M/N \text{ is open} & \text{ if and only if } U \subseteq M \text{ is open} \\ & \text{ if and only if for all } x \in U \text{ there exists } \lambda, x + M_\lambda \subseteq U \\ & \text{ if and only if for all } x \in \bar{U} \text{ there exists } \lambda \text{ with } \bar{x} + M'_\lambda \subseteq \bar{U} \\ & \text{ if and only if } \bar{U} \text{ is open in the linear topology} \end{aligned}$$

□

One can check that  $M/N$  is separated (i.e. the intersection of  $M_\lambda$ 's is 0) if and only if  $N \subseteq M$  is closed. Also check that the subspace topology on  $N$  coincides with the linear topology generated by  $\{N \cap M_\lambda\}_{\lambda \in \Lambda}$ .

Now, for every  $\lambda \in \Lambda$ , we have an exact sequence

$$0 \rightarrow N/N \cap M_\lambda \rightarrow M/M_\lambda \rightarrow (M/N)/M'_\lambda = M/(N + M_\lambda) \rightarrow 0$$

which is compatible with varying  $\lambda$ . Hence, we obtain the maps  $\hat{N} \rightarrow \hat{M} \rightarrow \widehat{M/N}$ . In fact, we get an exact sequence

$$0 \rightarrow \hat{N} \rightarrow \hat{M} \rightarrow \widehat{M/N}.$$

# Chapter 4

## Introduction to Homology

---